

FORM PTO-1390  
(REV 12-29-99)

U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE

ATTORNEY'S DOCKET NUMBER

**TRANSMITTAL LETTER TO THE UNITED STATES  
DESIGNATED/ELECTED OFFICE (DO/EO/US)  
CONCERNING A FILING UNDER 35 U.S.C. 371**

5368-US

U.S. APPLICATION NO. (If known, see 37 CFR 1.5)

09/582950

INTERNATIONAL APPLICATION NO.

PCT/US99/00419

INTERNATIONAL FILING DATE

08 January 1999

PRIORITY DATE CLAIMED

09 January 1998

**TITLE OF INVENTION**

SYNTHESIS OF COMBRETASTATIN A-4 PRODRUGS AND TRANS-ISOMERS THEREOF

**APPLICANT(S) FOR DO/EO/US**

G.R. PETTIT et al

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. ☒ This is a **FIRST** submission of items concerning a filing under 35 U.S.C. 371.
  2. ☐ This is a **SECOND** or **SUBSEQUENT** submission of items concerning a filing under 35 U.S.C. 371.
  3. ☐ This express request to begin national examination procedures (35 U.S.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39(1).
  4. ☒ A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date.
  5. ☒ A copy of the International Application as filed (35 U.S.C. 371(c)(2))
    - a. ☐ is transmitted herewith (required only if not transmitted by the International Bureau).
    - b. ☐ has been transmitted by the International Bureau.
    - c. ☒ is not required, as the application was filed in the United States Receiving Office (RO/US).
  6. ☐ A translation of the International Application into English (35 U.S.C. 371(c)(2)).
  7. ☐ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3))
    - a. ☐ are transmitted herewith (required only if not transmitted by the International Bureau).
    - b. ☐ have been transmitted by the International Bureau.
    - c. ☐ have not been made; however, the time limit for making such amendments has NOT expired.
    - d. ☐ have not been made and will not be made.
  8. ☐ A translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)).
  9. ☒ An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)).
  10. ☐ A translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).
- Items 11. to 16. below concern document(s) or information included:**
11. ☐ An Information Disclosure Statement under 37 CFR 1.97 and 1.98.
  12. ☐ An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
  13. ☐ A FIRST preliminary amendment.  
☐ A SECOND or SUBSEQUENT preliminary amendment.
  14. ☐ A substitute specification.
  15. ☐ A change of power of attorney and/or address letter.
  16. ☒ Other items or information: Copies of Verified Statement Claiming Small Entity Status - Independent Inventor as originally filed with US Provisional Application No. 60/071,070 filed January 9, 1998  
Certificate of Express Mail

U.S. APPLICATION NO. (if known, see 37 CFR 1.5)

09/582950

INTERNATIONAL APPLICATION NO.

PCT/US99/00419

ATTORNEY'S DOCKET NUMBER

5368-US

CALCULATIONS PTO USE ONLY

17. ☒ The following fees are submitted:**BASIC NATIONAL FEE (37 CFR 1.492 (a) (1) - (5)) :**

Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO and International Search Report not prepared by the EPO or JPO ..... \$970.00

International preliminary examination fee (37 CFR 1.482) not paid to USPTO but International Search Report prepared by the EPO or JPO ..... \$840.00

International preliminary examination fee (37 CFR 1.482) not paid to USPTO but international search fee (37 CFR 1.445(a)(2)) paid to USPTO ..... \$690.00

International preliminary examination fee paid to USPTO (37 CFR 1.482) but all claims did not satisfy provisions of PCT Article 33(1)-(4) ..... \$670.00

International preliminary examination fee paid to USPTO (37 CFR 1.482) and all claims satisfied provisions of PCT Article 33(1)-(4) ..... \$96.00

**ENTER APPROPRIATE BASIC FEE AMOUNT =**

\$ 96.00

Surcharge of \$130.00 for furnishing the oath or declaration later than ☐ 20 ☐ 30 months from the earliest claimed priority date (37 CFR 1.492(e)).

\$

CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE
Total claims	10 - 20 =	0	X \$18.00
Independent claims	3 - 3 =	0	X \$78.00
MULTIPLE DEPENDENT CLAIM(S) (if applicable)			+ \$260.00

\$ 0

\$ 0

\$

**TOTAL OF ABOVE CALCULATIONS =**

\$ 96.00

Reduction of 1/2 for filing by small entity, if applicable. A Small Entity Statement must also be filed (Note 37 CFR 1.9, 1.27, 1.28).

\$ 48.00

**SUBTOTAL =**

\$ 48.00

Processing fee of \$130.00 for furnishing the English translation later than ☐ 20 ☐ 30 months from the earliest claimed priority date (37 CFR 1.492(f)).

\$

**TOTAL NATIONAL FEE =**

\$ 48.00

Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31). \$40.00 per property

\$

**TOTAL FEES ENCLOSED =**

\$ 48.00

Amount to be refunded:

\$

charged:


\$

a. ☒ A check in the amount of \$ 48.00 to cover the above fees is enclosed.b. ☐ Please charge my Deposit Account No. \_\_\_\_\_ in the amount of \$ \_\_\_\_\_ to cover the above fees. A duplicate copy of this sheet is enclosed.c. ☒ The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. 13-5100. A duplicate copy of this sheet is enclosed.

**NOTE:** Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.

SEND ALL CORRESPONDENCE TO:

Richard R. Mybeck  
8010 East Morgan Trail, Suite 10  
Scottsdale, Arizona 85258-1234  
480.483,1285



SIGNATURE:

Richard R. Mybeck

NAME

17,886

REGISTRATION NUMBER

APPLICANT: GEORGE R. PETTIT et al. DOCKET NO. 5368  
SERIAL NO.:  
FILED:  
FOR: SYNTHESIS OF COMBRETASTATIN A-4 PRODRUG AND TRANS-ISOMER THEREOF

VERIFIED STATEMENT CLAIMING SMALL ENTITY  
STATUS - INDEPENDENT INVENTOR

As a below named inventor, I hereby declare that I qualify as an independent inventor as defined in 37 CFR 1.9(c) for purposes of paying reduced fees under section 41(a) and (b) of Title 35, United States Code, to the Patent and Trademark Office with regard to the invention entitled SYNTHESIS OF COMBRETASTATIN A-4 PRODRUG AND TRANS-ISOMER THEREOF described in the specification filed herewith.

I have not assigned, granted, conveyed or licensed and am under no obligation under contract or law to assign, grant, convey or license, any rights in the invention to any person who could not be classified as an independent inventor under 37 CFR 1.9(c) if that person had made the invention, or to any concern which would not qualify as a small business concern under 37 CFR 1.9(d) or a nonprofit organization under 37 CFR 1.9(e).

I acknowledge the duty to file, in this application, notification of any change in status resulting in loss of entitlement to small entity status prior to paying, or at the time of paying, the earliest of the issue fee or any maintenance fee due after the date on which status as a small entity is no longer appropriate. (37 CFR 1.28(b)).

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application and any patent issuing thereon to which this verified statement is directed.

Date 1/8/98

George R. Pettit  
GEORGE R. PETTIT

APPLICANT: GEORGE R. PETTIT et al. DOCKET NO. 5368  
SERIAL NO.:  
FILED:  
FOR: SYNTHESIS OF COMBRETASTATIN A-4 PRODRUG AND TRANS-ISOMER THEREOF

VERIFIED STATEMENT CLAIMING SMALL ENTITY  
STATUS - INDEPENDENT INVENTOR

As a below named inventor, I hereby declare that I qualify as an independent inventor as defined in 37 CFR 1.9(c) for purposes of paying reduced fees under section 41(a) and (b) of Title 35, United States Code, to the Patent and Trademark Office with regard to the invention entitled SYNTHESIS OF COMBRETASTATIN A-4 PRODRUG AND TRANS-ISOMER THEREOF described in the specification filed herewith.

I have not assigned, granted, conveyed or licensed and am under no obligation under contract or law to assign, grant, convey or license, any rights in the invention to any person who could not be classified as an independent inventor under 37 CFR 1.9(c) if that person had made the invention, or to any concern which would not qualify as a small business concern under 37 CFR 1.9(d) or a nonprofit organization under 37 CFR 1.9(e).

I acknowledge the duty to file, in this application, notification of any change in status resulting in loss of entitlement to small entity status prior to paying, or at the time of paying, the earliest of the issue fee or any maintenance fee due after the date on which status as a small entity is no longer appropriate. (37 CFR 1.28(b)).

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application and any patent issuing thereon to which this verified statement is directed.

Date January 7, 1998

Monte R. Rhodes  
MONTE R. RHODES

Synthesis of Combretastatin A-4 Prodrugs  
and Trans-isomers Thereof

5 The present invention relates generally to the field of compounds which may be useful in the treatment of one or more neoplastic diseases and more particularly, this invention relates to the synthesis of water soluble prodrugs of the antineoplastic compound denominated "combretastatin A-4."

10 This research was funded in part by Outstanding Investigator Grant CA44344-05-09 awarded by the National Cancer Institute, DHHS. The United States government may have certain rights to this invention.

The present invention relates to new and improved methods of synthesizing prodrugs and trans-isomers of the known antineoplastic compound denominated combretastatin A-4.

15 In addition, this invention teaches the chemistry of forming phosphate salts by reaction with the phenolic hydroxy group present in combretastatin A-4 and its analogs as described in U.S. Patents 4,940,726; 5,409,953; and 5,569,786.

20 combretastatin A-4 has poor water solubility distribution within biological systems. The elucidation and isolation of combretastatin A-4 is described in U.S. Patent No. 4,996,237 which issued to G.R. Pettit et al., on February 26, 1991, while early efforts to develop a combretastatin A-4 prodrug are described in U.S. Patent No. 5,561,122, which issued to G.R. Pettit on October 1, 1996. The general background information from each of those patents is incorporated herein by this reference thereto.

25 The potent cancer cell growth and tubulin assembly inhibitor combretastatin A-4 was originally isolated from the African tree *Combretum caffrum* (*Combretaceae*) circa 1985 and has been undergoing preclinical development. However, because of the very sparing aqueous solubility behavior of the phenol and its alkali metal salts, drug formulation attempts have given unsatisfactory results. The present invention represents a breakthrough in the

continuing effort to synthesize practical water soluble prodrugs based on combretastatin A-4 and is a significant advance over the early efforts described in U.S. Patent No. 5,561,122, *supra*.

5 In certain African Zulu traditional practices, the root bark of *Combretum caffrum* (*C. salicyolium*, *combretaceae* Family) is used as a charm for causing harm to an enemy. Combretastatin A-4, **1a**, a *Z*-stilbene, isolated (Pettit *et al.*, 1989, Isolation and structure of the strong cell-growth and tubulin inhibitor combretastatin A-4, *Experimentia*, 45, 209) from this tree has been shown to be a potent cancer cell growth inhibitor (El-Zayat *et al.*, 1993, *In vitro* evaluation of the antineoplastic activity of combretastatin A-4, a natural product from *Combretum caffrum* (arid shrub), *Anti-Cancer Drugs*, 4, 19); cancer anti-agent (Dark *et al.*, 1997, combretastatin A-4, an agent that displays potent and selective toxicity towards tumour vasculature); and a tubulin assembly inhibitor (Lin *et al.*, 1989, Antimitotic natural products combretastatin A-4 and combretastatin A-2: studies on the mechanism of their inhibition of the binding of colchicine to tubulin, *Biochemistry*, 28, 6984). Combretastatin a-4 is now in clinical development.

10 Although combretastatin A-4 has a phenol hydroxyl group and has demonstrated considerable promise as a unique anticancer agent, its development has been inhibited by its extremely poor solubility in water. A series of water-soluble derivatives have been recently reported (Brown *et al.*, 1995, Synthesis of water-soluble sugar derivatives of combretastatin A-4, *Journal of the Chemical Society, Perkin Transactions I*, 577; Woods *et al.*, 1995, The interaction with tubulin of a series of stilbenes based on combretastatin A-4, *British Journal of Cancer*, 71, 705; and, Bedford *et al.*, 1996, Synthesis of Water-Soluble Prodrugs of the Cytotoxic Agent combretastatin A-4, *Bioorganic & Medicinal Chemistry Letters*, 6, 157) where the water-soluble phosphate sodium salt, **1h**, (Pettit *et al.*, 1995, Antineoplastic agents 322, Synthesis of combretastatin A-4 prodrugs, *Anti-Cancer Drug Design*, 10, 299; and, Pettit, U.S.

Patent No. 5,561,122) proved to be the most attractive. However, the phosphorylation sequence employing *bis*(2,2,2-trichloroethyl)phosphorochloridate, subsequent reduction (Zn, CH<sub>3</sub>CO<sub>2</sub>H), and isolation by ion-exchange chromatography was not well suited to producing prodrug **1h** on a large scale and as such, posed a significant economic obstacle to its ultimate commercialization.

While prodrug options are steadily increasing and include poly(ethylene glycol) esters (Greenwald *et al.*, 1996, Drug delivery systems: water soluble taxol 2'-poly(ethylene glycol) ester prodrugs-design and *in vivo* effectiveness, *The Journal of Medicinal Chemistry*, 39, 424); quaternary salts (Lackey *et al.*, 1996, Water soluble inhibitors of topoisomerase I: quaternary salt derivatives of camptothecin, *The Journal of Medicinal Chemistry*, 39, 713); sulfonate salts (Hejchman *et al.*, 1995, Synthesis and cytotoxicity of water-soluble ambrosin prodrug candidates, *The Journal of Medicinal Chemistry*, 38, 3407); urethans (Izawa *et al.*, 1995, Design and synthesis of an antitumor prodrug released by the reaction with sulfhydryl compounds, *Bioorganic & Medicinal Chemistry Letters*, 5, 593); biodegradable polymers (Gombotz *et al.*, 1995, Biodegradable polymers for protein and peptide drug delivery, *Bioconjugate Chemistry*, 6, 332); and, a variety of other methods (Jungheim *et al.*, 1994, Design of antitumor prodrugs: substrates for antibody targeted enzymes, *Chemical Reviews*, 94, 1553), the present disclosure is focused on the previously introduced sodium phosphate derivative **1h** (which is equivalent to the sodium phosphate derivative **3d**, produced by the methods set forth herein, see below) of combretastatin A-4. From evidence at hand phosphate **1h** (as well as **3d**) is presumed to be appropriately dephosphorylated by serum phosphatases and then transported intracellularly.

Furthermore, the phosphate prodrug approach has proved useful with substances as diverse as Etoposide (Saulnier *et al.*, 1994, Synthesis of etoposide phosphate, BMY-40481: a water-soluble clinically active prodrug of etoposide,

*Bioorganic & Medicinal Chemistry Letters*, 4, 2567); Taxol (Mamber *et al.*, 1995, Tubulin polymerization by paclitaxel (Taxol) phosphate prodrugs after metabolic activation with alkaline phosphatase, *The Journal of Pharmacology and Experimental Therapeutics*, 374, 877; Ueda *et al.*, 1995, Novel, water-soluble phosphate derivatives of 2'-ethoxycarbonylpaclitaxel as potential prodrugs of paclitaxel: synthesis and antitumor evaluation, *Bioorganic & Medicinal Chemistry Letters*, 5, 247; and Ueda, *et al.*, 1993, Novel water soluble phosphate prodrugs of Taxol possessing *in vivo* antitumor activity, *Bioorganic & Medicinal Chemistry Letters*, 3, 1761); and tyrosine-containing peptides (Chao *et al.*, 1993, N,N-Diisopropyl-bis[2-(trimethylsilyl)ethyl]phosphoramidite, An attractive phosphorylating agent compatible with the Fmoc/t-butyl strategy for the synthesis of phosphotyrosine containing peptides, *Tetrahedron Letters*, 34, 3377). As shown herein below for prodrug **1h** (also, **3d**), phosphate ester salts are generally cleaved *in vivo* (Bundgaard (ed.), 1985, Design of prodrugs, pp. 1-92, Elsevier: New York) and are stable enough to usually allow formulation into solutions with practical shelf lives (Flynn *et al.*, 1970, Factors influencing solvolysis of corticosteroid-21-phosphate esters, *Journal of Pharmaceutical Sciences*, 59, 1433). To improve the synthetic scaleup practicality for the sodium phosphate ester prodrug **1h** (**3d**) of combretastatin A-4, three new phosphate esters have been synthesized which were easily transformed into the water-soluble prodrug **3d** as is herein described.

The aggressive behavior of a broad spectrum of human cancer types such as breast carcinoma is related to the facility for rapid tumor angiogenesis. Neovascularization represents a very attractive target for new tumor antiangiogenesis-type anticancer drugs. Combretastatin A-4 **1a** was found to show potent antiangiogenesis activity and the derived sodium phosphate prodrug **1h** (**3d**) is a powerful *in vivo* inhibitor of tumor vascularization. Currently prodrug **1h** is in clinical development. A practical synthetic procedure for providing the pure drug in quantity for clinical trials has




continued to be an important research objective. As described, the present invention is predicated upon the discovery of a considerably improved sequence utilizing *in situ* prepared dibenzyl chlorophosphite for phosphorylating phenol **1a**, cleavage of the benzyl ester groups by trimethyliodosilane and treatment of the product with sodium methoxide to afford prodrug **1h** (herein **3d**) in good yield. However, the product was found to be accompanied by an isomeric substance that was first revealed by the appearance of some opalescence in aqueous solutions of prodrug **3d**. The side product was believed to be the corresponding *trans*-stilbene **4c** and was eliminated by extraction and fractional recrystallization. The purity of prodrug **3d** was readily assigned by an ion-pair HPLC analysis using a phosphate buffer.


This further study was focused on confirming the structure of the persistent by-product (assumed to be **4c**) by synthesis. In addition, the evaluation of combretastatin A-4 prodrugs was extended by preparing an extended series of phosphate metal and ammonium cation derivatives of the phosphoric acid precursor of prodrug **3d**. Synthesis of the *trans*-stilbene phosphate **4c** was also considered potentially useful for SAR purposes in view of the chemopreventative activities of certain *trans*-stilbene phenols such as resveratrol.

Combretastatin A-4 is essentially insoluble in water. This characteristic has significantly interfered with accomplishing the necessary formulations of pharmaceutical preparations of this compound for use in pre-clinical development. Hence, derivatives of the combretastatin A-4, **1a**, 3' phenol group were prepared for evaluation as possible water soluble prodrugs. As noted in U.S. Patent No. 5,561,122, the sodium salt **1b**, potassium salt **1c** and hemisuccinic acid ester **1d** derivatives of phenol **1a** were essentially insoluble in water. Indeed, these substances regenerated combretastatin A-4 upon reaction with water. A series of other simple derivatives proved unsatisfactory in terms of water solubility or stability or both. The most soluble derivatives

evaluated included the ammonium **1f**, potassium **1g** and sodium **1h** phosphate salts where the latter two proved most stable and suitable. Both the sodium and other phosphate salt derivatives of combretastatin A-4 described herein were also found to exhibit the requisite biological properties necessary for a useful prodrug.



- 1a**, R = H  
**b**, R = Na  
**c**, R = K  
**d**, R = COCH<sub>2</sub>CH<sub>2</sub>CO<sub>2</sub>H




**f**, R = ammonium phosphate salt

**g**, R = potassium phosphate salt


**h**, R = sodium phosphate salt, e.g.

Combretastatin A-4 **1a**, as the phosphate ester prodrug **1h** (and, hereinafter **3d**) is a potent antineoplastic and antiangiogenesis substance and is in clinical development. For the purpose of improving the phosphorylation synthetic sequence from combretastatin A-4, new routes were investigated. The phosphorylation step was found to be considerably improved using *in-situ* generated dibenzyl chlorophosphite. Cleavage of the benzyl esters employing a trimethylchlorosilane/sodium iodide procedure, followed by treatment with sodium methoxide, led to the water-soluble prodrug **3d** in high yield.




As shown more particularly above for prodrug **3d**, phosphate ester salts are generally cleaved *in vivo* (Bundgaard, *supra*). To improve the synthetic scaleup practicality for the sodium phosphate ester prodrug of combretastatin A-4, three new phosphate esters **3a**, **3b**, and **3c** have been synthesized the latter of which was easily transformed into the water-soluble prodrug **3d** as is described in more detail below.

The (E)-stilbene isomer **4c** (see below) of the natural (Z)-combretastatin A-4 prodrug **3d** was also efficiently prepared from (E)-combretastatin A-4 **1a** by an analogous reaction sequence employing phosphorylation (dibenzyl chlorophosphite), cleavage (trimethyliodosilane) of the benzyl ester and reaction of the resulting phosphoric acid with sodium methoxide. The sodium phosphate product **4c** was found to be an important side-product, presumably from iodine-catalyzed isomerization, when this synthetic route was used to obtain the combretastatin A-4 prodrug **3d**.




**1a**, R = OH  
**3d**, R = OPO<sub>3</sub>H(Na)



**4a**, R = OH  
**b**, R = OPO<sub>3</sub>Bn<sub>2</sub>  
**c**, R = OPO<sub>3</sub>H(Na)

The phosphoric acid precursor of prodrug **3d** derived from (Z)-combretastatin A-4 **1a** was also converted to a series of metal cation and ammonium cation salts to evaluate effects on human cancer cell growth, antimicrobial activities and solubility behavior. Thus, several new combretastatin A-4 prodrug modifications **5a-s** (see below) have been prepared and found to exhibit inhibitory action against various bacteria and opportunistic fungi.




where X=H(Z) (monovalent)  
 or  
 X=Z (divalent)

**5a** Z = Ca<sup>2+</sup>  
**b** Z = Cs<sup>2+</sup>  
**c** Z = Li<sup>+</sup>  
**d** Z = Mg<sup>2+</sup>  
**e** Z = Mn<sup>2+</sup>  
**f** Z = Zn<sup>2+</sup>

**g** Z = imidazole  
**h** Z = morpholine  
**i** Z = piperazine  
**j** Z = piperidine  
**k** Z = pyrazole  
**l** Z = pyridine

**m** Z = adenosine  
**n** Z = cinchonine  
**o** Z = glucosamine  
**p** Z = quinine  
**q** Z = quinidine  
**r** Z = tetracycline  
**s** Z = verapamil



verapamil

Accordingly, the prime object of the subject invention is to prepare prodrugs of combretastatin A-4 which are both water soluble and stable, as well as to improve the means of synthesizing such compounds.

These and still further objects as shall hereinafter appear are readily fulfilled by the present invention in a remarkably unexpected manner as will be readily discerned from the following detailed description of an exemplary embodiment thereof especially when read in conjunction with the structures set forth above and the accompanying drawing.

FIG. 1 comprises high performance liquid chromatography charts for a) mixture of (Z)- and (E)-combretastatin A-4 prodrugs b) combretastatin A-4 prodrug, 100% pure by HPLC.

The process by which combretastatin may be synthesized is well known in the art. *See*, Pettit et al., Isolation and structure of the strong cell growth and tubulin inhibitor combretastatin A-4, *Experimentia*, 209 (1989), *supra*. Combretastatin A-4, synthesized by that method disclosed in U.S. Patent No. 4,996,237 ("the '237 patent"), was used throughout this process.

Combretastatin A-4 (**1a**) was synthesized as previously described by Pettit *et al*, in 1995. Diisopropylethylamine (98%), lithium hydroxide monohydrate, cesium hydroxide, zinc acetate dihydrate, quinine monohydrate, piperidine, pyrazole, and sodium methoxide (95%), dibenzyl phosphite, 4-dimethylaminopyridine (DMAP), 1*H*-Tetrazole (98%), 85% 3-chloroperoxybenzoic acid (*m*-CPBA), chlorotrimethylsilane (98%), trifluoroacetic acid (99%), triethylamine, and sodium methoxide (95%) were obtained from Sigma-Aldrich Chemical Company. The *tert*-butanol was obtained from Acros Organics. Calcium acetate, magnesium acetate tetrahydrate, manganese acetate and other reaction and isolation reagents were obtained from the Baker Chemical Company. Tetrahydrofuran was distilled from sodium/benzophenone. Imidazole was recrystallized from *n*-hexane, and piperazine was recrystallized from ethanol and dried *in vacuo*. The morpholine was dried with KOH and then fractionally distilled; pyridine was dried over CaH<sub>2</sub> and fractionally distilled, then stored over KOH. Adenosine was obtained from Nutritional Biochemicals Corporation; cinchonine, from Mateson, Cole & Bell and was recrystallized from ethanol prior to use; D-glucosamine was from Nutritional Biochemicals Corporation; quinidine, was from J. T. Baker; tetracycline, from Mateson, Cole & Bell; and verapamil, from the Alexis Corporation. "Ether", refers to diethyl ether. All solvents were redistilled prior to use. The 1.0 M solutions of the metal salts were in distilled water, and the 1.0 M solutions of the amines were in dry methanol. The di-*tert*-butyloxy(N,N-diethylamido)phosphine was prepared as previously described by Johns *et al* (1988, Di-*tert*-butyl N,N-Diethylphosphoramidite, A New Phosphitylating Agent for the Efficient Phosphorylation of Alcohols, *Synthesis*, pp. 142-144).

Reactions were monitored by thin-layer chromatography using ANALTECH silica gel GHLF Uniplates visualized under long-wave and short-wave UV irradiation. Solvent extracts of aqueous solutions were dried over anhydrous sodium sulfate. Where appropriate, the crude products were

separated by flash column chromatography (230-400 mesh ASTM) from E. Merck.

Melting points were measured with an ELECTROTHERMAL digital melting point apparatus, model IA9200, and are uncorrected. The IR spectra were obtained using a NICOLET FTIR Model MX-1 instrument. EIMS data was recorded with a MAT 312 mass spectrometer, and high-resolution FAB spectra were obtained with a KRATOS MS-50 mass spectrometer (Midwest Center for Mass Spectrometry, University of Nebraska, Lincoln, NE). All  $^1\text{H}$ - and  $^{13}\text{C}$ -NMR spectra were obtained using a VARIAN GEMINI 300 MHz instrument with  $\text{CDCl}_3$  (TMS internal reference) as solvent unless otherwise noted. The  $^{31}\text{P}$ -NMR spectra were obtained in  $\text{CDCl}_3$  with 85%  $\text{H}_3\text{PO}_4$  as an external standard employing a UNITY 500 MHz instrument. Elemental analyses were determined by Galbraith Laboratories, Inc., Knoxville, TN.

#### HPLC Separations and Analysis

HPLC analyses were performed on a HEWLETT-PACKARD HP 1050 Series 3D CHEMSTATION, Rev. A.03.01, liquid chromatograph. All solvents used were HPLC grade and filtered before use. HPLC separation and analysis procedures were based on aqueous buffered ion-pair reverse-phase gradient elution at  $33^\circ\text{C}$ . Flow rate: 1.2 ml/min. Column: MERK LICHROSPHERE 100 RP 18 ( $5\mu\text{m}$ ), 250 x 4.0 mm (LICHROART 250-4). Eluents: Eluent A. 5mM tetrabutylammonium hydrogen sulfate, 5 mM  $\text{H}_3\text{PO}_4$ , 5 mM  $\text{KH}_2\text{PO}_4$ . Eluent B. 75%  $\text{CH}_3\text{CN}$ . Gradient: a) Hold 4 minutes at 65% A, 35% B, b) 8 minutes linear gradient, 65% A to 10% A, c) Hold 3 minutes. Detector: 310 nm UV.

#### *di-tert-Butyloxy(N,N-diisopropylamide)phosphine (2a)*

A solution of *tert*-butanol (3.7 g, 4.7 ml, 50 mmol, 2.0 equiv) and triethylamine (5.6 g, 7.7 ml, 55 mmol, 2.2 equiv) in dry ether (20 ml) was added to dichloro-N,N-diisopropylphosphine (5.0 g, 25 mmol) in dry ether (5 ml) while maintaining the solution temperature below  $0^\circ\text{C}$ . Upon completion

of addition (about 15 minutes), the solution was allowed to warm to room temperature and stirred for a further 3 hours. A solution of 5% aqueous sodium bicarbonate (10 ml) was added, the aqueous phase was removed, and the ethereal solution was washed with 5% aq. sodium bicarbonate (2 x 10 ml), saturated aq. NaCl (25 ml), dried, filtered, and solvent removed to give a clear residual oil. Purification by vacuum distillation gave the product as a clear oil (4.3 g, 63%). b.p. 56-57°C/0.05 mm Hg; <sup>1</sup>H-NMR (300 MHz) δ 1.17 (d, *J*=6.0 Hz, CH(CH<sub>3</sub>)<sub>2</sub>), 1.35 (s, 18H, 2 x (CH<sub>3</sub>)<sub>3</sub>C), 3.53-3.68 (m, 2 x (CH<sub>3</sub>)<sub>2</sub>CH); <sup>13</sup>C-NMR (125 MHz) δ 24.21 (d, *J*<sub>PC</sub>=7.73 Hz), 31.06 (d, *J*<sub>PC</sub>=9.75 Hz), 43.09 (d, *J*<sub>PC</sub>=13.88 Hz), 74.50 (d, *J*<sub>PC</sub>=9.75 Hz); <sup>31</sup>P-NMR (202 MHz) δ 130.0 (s).

*di-tert-Butyloxy(N,N-diethylamido)phosphine (2b)*

Phosphine **2b** was as previously described (Johns *et al*, 1988, *supra*): b.p. 49-51°C/0.005 mm Hg (55%yield); <sup>1</sup>H-NMR (300 MHz) δ 1.06 (t, 6H, *J*<sub>HH</sub>=7.8 Hz), 1.33 (s, 18H, C(CH<sub>3</sub>)<sub>3</sub>), 3.03 (dq, 4H, *J*<sub>HH</sub>=7.8 Hz, *J*<sub>PH</sub>=9.0 Hz, PCH<sub>2</sub>CH<sub>3</sub>); <sup>13</sup>C-NMR (75 MHz) δ 14.84 (d, *J*<sub>PC</sub>=3.9 Hz), 30.88 (d, *J*<sub>PC</sub>=9.75 Hz), 37.45 (d, *J*<sub>PC</sub>=21.3 Hz), 74.40 (d, *J*<sub>PC</sub>=11.63 Hz).

*Bis-[2-(trimethylsilyl)ethoxy]-N,N-diisopropylamidophosphine (2c)*

A solution of 2-(trimethylsilyl)ethanol (1.2 g, 9.9 mmol, 2 equiv) and triethylamine (1.5 ml, 10.9 mmol, 2.2 equiv) in dry ether (10 ml) was added to dichloro-N,N-diisopropylamidophosphine (1.0 g, 4.9 mmol) in dry ether (10 ml) and the temperature was maintained below 0°C. Upon completion of addition (*ca* 30 min), the mixture was allowed to warm to room temperature and stirring continued for 3 hours. After adding a solution of 5% aq. NaHCO<sub>3</sub> (10 ml) the aqueous phase was washed with 5% aq. NaHCO<sub>3</sub> (2 x 10 ml), saturated aq NaCl (10 ml), dried and filtered. Removal of solvent *in vacuo* afforded a clear oil (2.2 g, 78%). Purification by vacuum distillation gave the product as a clear oil. b.p. 62-64°C/0.005 mm; IR (neat) ν<sub>max</sub> 2960 (s), 2897 (m), 1462 (w), 1396 (m), 1363 (m), 1249 (s), 1186 (s), 1126 (m) 1045 (s), 972 (s), 837 (s), 744 (s), 694 (s), 663 (w) cm<sup>-1</sup>; <sup>1</sup>H-NMR (300 MHz) δ 0.00 (s, 18H), 1.02 (m, 4H) 1.16



(d,  $J=6.6$  Hz, 12H), 3.52-3.77 (m, 6H);  $^{13}\text{C}$ -NMR (75 MHz)  $\delta$  -1.49, 20.00, 24.41, 24.51, 42.47, 42.63, 60.43, 60.67;  $^{31}\text{P}$ -NMR (202 MHz)  $\delta$  143.50.

*3'-O-Bis(benzyl)phosphorylcombretastatin A-4 (3a)*

A flame-dried, 3-neck flask containing a teflon stirbar was fitted with a septa, thermometer, and argon inlet. The phenol (**1a**, 20.0 g, 63.2 mmol) was dissolved in acetonitrile (200 ml) with stirring and then cooled to  $-25^{\circ}\text{C}$ . Carbon tetrachloride (30.5 ml, 316 mmol, 5 equiv) was added and the solution stirred for 5 minutes. Diisopropylethylamine (23.13 ml, 133 mmol, 2.1 equiv) was added *via* syringe, followed by DMAP (772 mg, 6.32 mmol, 0.1 equiv). One minute later dropwise (slow) addition of dibenzyl phosphite (20.33 ml, 92 mmol, 1.45 equiv) was begun at such a rate that the reaction temperature was kept below  $-10^{\circ}\text{C}$ . When the reaction was complete (1 hr by TLC analysis) 0.5 M aqueous  $\text{KH}_2\text{PO}_4$  was added (50 ml) and the solution allowed to warm to room temperature. After extraction with ethyl acetate (3 x 100 ml), the combined solvent extract was washed with water (100 ml), saturated NaCl (100 ml) and dried. Filtration and removal of solvent under reduced pressure gave a faint yellow oil. The oily residue was chromatographed on a column of silica gel (hexane-ethyl acetate, 3:2) to give a clear oil which crystallized from ethyl acetate-hexane to give pure, colorless needles (36.0 g; 98%): m.p.  $73^{\circ}\text{C}$ ;  $R_f$  0.26 (hexane-ethyl acetate, 3:2); EIMS  $m/z$  576 (100,  $\text{M}^+$ ), 561 (8), 485 (2), 394 (2), 378 (3), 363 (3), 252 (5), 181 (3); IR (neat)  $\nu_{\text{max}}$  3005, 2939, 2837, 1579, 1512, 1454, 1425, 1280, 1240, 1122, 1010, 891, 740, 698  $\text{cm}^{-1}$ ;  $^1\text{H}$ -NMR (300 MHz)  $\delta$  3.67 (s, 6H, 2 x  $\text{OCH}_3$ ), 3.77 (s, 3H,  $\text{OCH}_3$ ), 3.80 (s, 3H,  $\text{OCH}_3$ ), 5.12 (s, 2H,  $\text{CH}_2\text{-Ar}$ ), 5.14 (s, 2H,  $\text{CH}_2\text{-Ar}$ ), 6.40 (d,  $J=12.0$  Hz, 1H), 6.45 (d,  $J=12.0$  Hz, 1H), 6.48 (s, 2H), 6.78 (d,  $J=9.0$  Hz, 1H), 7.06 (dd,  $J=9.0, 1.8$  Hz, 1H), 7.15 (d,  $J=1.8$  Hz, 1H), 7.31 (s, 1OH, Ar-H);  $^{13}\text{C}$ -NMR (75 MHz)  $\delta$  152.97, 139.32, 139.22, 135.67, 132.46, 130.19, 129.67, 128.53, 127.88, 126.56, 122.21, 112.27, 105.93, 69.79, 69.72, 60.86, 60.40, 55.92, 21.05, 14.22;  $^{31}\text{P}$ -NMR

(202 MHz)  $\delta$  -5.29; Anal. Calcd. for  $C_{32}H_{33}O_8P$ : C, 66.67; H, 5.77. Found: C, 66.96; H, 6.05.

*3'-O-Bis(tert-butyl)phosphorylcombretastatin A-4 (3b)*

*Method A:* 1H-Tetrazole (3.73 g, 53 mmol, 3.67 equiv) was added in one portion to a stirred solution of combretastatin A-4 (**1a**, 5.00 g, 16 mmol, 1.0 equiv) and di-tert-butyloxy-N,N-diethylamidophosphine (5.19 g, 21 mmol, 1.3 equiv) in dry tetrahydrofuran (15 ml). The solution was stirred under argon for 20 minutes at room temperature and cooled to -70°C (CO<sub>2</sub>/acetone). Next a solution of 85% *m*-CPBA (3.22 g, 19.0 mmol) in dry dichloromethane (10 ml) was added (rapidly) such that the reaction mixture temperature was kept below -30°C. After stirring for 5 min at room temperature, 10% aqueous sodium bicarbonate (15 ml) was added. The solution was stirred for another 10 minutes and extracted with ether (70 ml). The ethereal phase was washed with 10% aq. sodium bicarbonate (2 x 20 ml), 1N NaOH (6 x 40 ml), dried and filtered. Evaporation of solvent under reduced pressure gave a yellow oil which was purified by flash column chromatography (silica gel; 3:2 hexane-ethyl acetate) to give the title compound as a clear oil (1.15 g; 75%): *R*<sub>f</sub> 0.25 (hexane-ethyl acetate, 3:2); EIMS *m/z* 508 (*M*<sup>+</sup>, 22), 452 (5), 437 (7), 396 (100), 381 (17), 364 (4), 349 (3), 316 (5); IR (neat)  $\nu_{\max}$  3462, 2980, 2937, 2837, 2243, 1579, 1512, 1462, 1425, 1371, 1327, 1276, 1174, 1128, 999, 920, 868, 731, 646 cm<sup>-1</sup>; <sup>1</sup>H-NMR (300 MHz)  $\delta$  1.47 (s, 18H, 2 x C(CH<sub>3</sub>)<sub>3</sub>), 3.69 (s, 6H, 2 x OCH<sub>3</sub>), 3.82 (s, 3H, OCH<sub>3</sub>), 3.83 (s, 3H, OCH<sub>3</sub>), 6.42 (d, *J*=12 Hz, 1H), 6.47 (d, *J*=12 Hz, 1H), 6.51 (s, 2H, 2,6-H), 6.79 (d, *J*=8.5 Hz, 1H), 7.03 (dd, *J*=8.5, 1.8 Hz, 1H), 7.31 (d, *J*=1.8 Hz, 1H); <sup>13</sup>C-NMR (75 MHz)  $\delta$  152.94, 132.62, 129.85, 129.25, 129.02, 125.63, 121.80, 112.14, 105.93, 99.28, 83.49, 60.89, 55.92, 29.80, 29.72; <sup>31</sup>P-NMR (202 MHz, CDCl<sub>3</sub>/ 85% H<sub>3</sub>PO<sub>4</sub>)  $\delta$  -14.47; Anal. calcd. for  $C_{26}H_{37}O_8P$ : C, 61.40%; H, 7.33%. Found: C, 60.92; H, 7.67.

*Method B:* The same procedure as in Method A above was followed except di-tert-butyloxy-N,N-diisopropylamidophosphine was used as the phosphorylating agent. Yield: 78%.

*3'-O-Bis(trimethylsilylethoxy)phosphorylcombretastatin A-4 (3c)*

The same procedure as in Method A above was used, with *bis*-[2-(trimethylsilyl) ethoxy-N,N-diisopropylamidophosphine (**2c**), giving the product as a colorless oil (8.67g, 91%):  $R_f$  0.26(hexane-ethyl acetate, 3:1); EIMS  $m/z$  596 ( $M^+$ , 90), 568 (20), 540 (21), 525 (38), 496 (10), 468 (40), 453 (13), 396 (22), 283 (10), 252 (11), 211 (13), 147 (15), 73 (100); IR (neat)  $\nu_{max}$  2999, 2955, 2901, 2837, 1579, 1512, 1464, 1525, 1276, 1249, 1217, 1180, 1128, 991, 856, 767, 696  $cm^{-1}$ ;  $^1H$ -NMR (500 MHz)  $\delta$  0.00 (s, 18 H), 1.04 (m, 4H,  $-CH_2Si-$ ), 3.63 (s, 6H, 2 x  $OCH_3$ ), 3.77 (s, 6H, 2 x  $OCH_3$ ), 4.17 (m, 4H,  $-OCH_2CH_2-$ ), 6.09 (s, 2H), 6.42 (d,  $J=12.5$  Hz, 1H), 6.47 (d,  $J=12.5$  Hz, 1H), 6.77 (d,  $J=8.5$  Hz, 1H), 7.03 (dd,  $J=8.5, 1.0$  Hz, 1H), 7.21 (d,  $J=1.0$  Hz, 1h);  $^{13}C$ -NMR (75 MHz)  $\delta$  152.77, 149.69, 139.54, 137.05, 132.24, 129.94, 129.37, 128.41, 126.05, 121.78, 112.08, 105.76, 66.72, 64.00, 60.59, 55.70, 19.51, 19.20, -1.72;  $^{31}P$ -NMR (202 MHz)  $\delta$  -5.54.

*Sodium (Z)-combretastatin A-4 3'-O-phosphate (3d)*

*Procedure A:* To a solution of the benzyl ester (**3c**, 20.5 g, 35.6 mmol) and sodium iodide (10.7 g, 71 mmol, 2 equiv) in dry acetonitrile (100 ml, in a flame-dried flask under argon) was slowly added chlorotrimethylsilane (9.02 ml, 71 mmol, 2 equiv) with vigorous stirring. The reaction mixture was stirred for 20 minutes at which point TLC analysis (hexane-ethyl acetate, 3:2) showed no starting material. Water (just enough to dissolve the salts) was added and the straw color removed by the addition of 10% aq.sodium thiosulfate (2 ml). The organic phase was separated and the aqueous phase extracted with ethyl acetate (4 x 50 ml). The combined extract was concentrated *in vacuo* to give a pale yellow foam. The foam was dissolved in dry methanol (100 ml) and sodium methoxide (95%, 4.1 g, 71 mmol, 2 equiv) was added in one portion and the

5 solution stirred for 9 hours. The methanol was removed *in vacuo* and the solid recrystallized from water-acetone and methanol-acetone to give colorless crystals (14 g, 90%): m.p. 190-195°C (decomp);  $R_f$  0.64 (n-butanol-methanol-water-ammonia, 4:3:2:1); UV  $\lambda_{max}$  (log  $\epsilon$ ) in CH<sub>3</sub>OH, 200 (4.54), 278 (4.25) nm (4.54), 196 (4.48) nm; IR (KBr)  $\nu_{max}$  2941 m, 2839s, 2361m (broad), 1581s, 1514s, 1431m, 1240m, 1124s, 983m, cm<sup>-1</sup>; <sup>1</sup>H-NMR (500 MHz, D<sub>2</sub>O)  $\delta$  3.50 (s, 6H, 3,5-OCH<sub>3</sub>), 3.57 (s, 3H, 4'-OCH<sub>3</sub>), 3.68 (s, 3H, 4-OCH<sub>3</sub>), 6.26 (d, 1H,  $J=12.5$  Hz, H-1a'), 6.42 (s, 2H, H-2,6), 6.44 (d, 1H,  $J=12.5$  Hz, H-1a), 6.63 (d, 1H,  $J=8.5$  Hz, H-5'), 6.71 (dd, 1H,  $J=2, 8.5$  Hz, H-6'), 7.19 (d, 1H,  $J=2$  Hz, H-2'); <sup>13</sup>C-NMR (125 MHz, D<sub>2</sub>O)  $\delta$  153.17, 150.50, 143.40, 136.90, 134.65, 131.28, 130.97, 129.93, 124.69, 122.64, 113.58, 107.54, 61.97, 57.10, 56.90; HRFAB  $m/z$  441.06942 [M + Na]<sup>+</sup>, 419 [M + H], 396 [M-Na + H]<sup>+</sup>. calc. 441.06912 for C<sub>18</sub>H<sub>20</sub>O<sub>8</sub>PNa<sub>2</sub>, anal. Na, 5.04%, H<sub>2</sub>O 6.26% calc. for C<sub>18</sub>H<sub>20</sub>O<sub>8</sub>PNa·1½ H<sub>2</sub>O hydrate, Na 5.16%, H<sub>2</sub>O 6.07%.

15 Sodium phosphate **3d** prepared by Procedure A was identical (TLC and spectra) with an earlier specimen (**1h**, Pettit *et al.*, 1995, *supra*; and U.S. Patent No. 5,561,122) obtained by the initial 3'-O-bis(2,2,2-trichloroethyl)phosphoryl route and exhibited solubility in distilled water of >20 mg/ml at 25°C.

20 *Procedure B*: A mixture of the di-*tert*-butylphosphoryl ester **3b** (7.25 g) and trifluoroacetic acid (25 ml) in dry dichloromethane (25 ml) was stirred at ambient temperature for 45 minutes (cleavage complete by TLC analysis hexane-acetone, 3:1). Water (10 ml) was added and solvent evaporated. Residual TFA was removed azeotropically with toluene (3 x 20 ml), and the resulting oily residue was dried under vacuum (0.005 mm/ Hg) for 1 hour. The oil was dissolved in dry methanol (2 ml) and sodium methoxide (1.60 g, 28 mmol, 2.0 equiv) was added. The solution was stirred under argon for 14 hours, solvent was removed *in vacuo* and the remaining solid dried. The residue was dissolved in water (10 ml) and the solution filtered through a fine porosity sintered glass funnel. Concentration of the filtrate gave a white solid which

recrystallized from water-acetone to give a colorless powder (5.8 g, 92%) identical with the sample of prodrug **3d** prepared by Procedure A.

*Procedure C:* Tetrabutylammonium fluoride (8.38 ml, 1.0 M in THF; 8.38 mmol) was slowly added employing a syringe to a stirred (under nitrogen) solution of di-silyl ethoxyphosphoryl ester **3c** (5.0 g, 8.38 mmol) in 15 ml of anhydrous tetrahydrofuran. After 30 minutes, TLC analysis (hexane-ethyl acetate, 3:2) indicated the reaction was complete. Ice (5 g) was added, followed by ether (50 ml). The organic solvent was separated, washed with water (3 x 50 ml), dried, filtered and solvent removed *in vacuo* to yield a straw colored gum. The product was dissolved in methanol (5 ml) and sodium methoxide (0.90 g, 16.8 mmol, 2 equiv) was added. The solution was stirred under argon for 12 hours, solvent was removed under reduced pressure and the residual solid was recrystallized from water-acetone to give a colorless powder (**3d**, 3.60, 98%) identical with the product from Procedure A.

After completing the first synthesis of sodium combretastatin A-4, 3'-O-phosphate **3d** a number of other phosphorylation approaches were explored toward the objective of establishing a procedure more readily adapted to large scale synthesis. Selection of an appropriate method for phosphorylation of combretastatin A-4 **1a** was also limited by available methods for removal of the eventual phosphate protecting groups. An alternative dibenzylphosphite-carbon tetrachloride procedure proved very practical for obtaining phosphate ester **3a** in high yield (95%). Catalytic hydrogenolysis of the benzyl oxygen bonds under varied conditions gave a concomitant reduction of the olefin bridge. Other benzyl oxygen cleavage methods investigated included the use of triphenylcarbonium tetrafluoroborate, arylthiotrimethylsilane-zinciodide, acidic alumina in a microwave and the Lewis acid SnCl<sub>4</sub> (Mukai *et al.*, 1996, Studies on the total synthesis of antitumor styryllactones: stereoselective total syntheses of (+)-goniofufurone, (+)-goniobutenolide A, and (-)-goniobutenolide B, *Tetrahedron*, 52, 6547). While all cleaved the benzyl group, they appeared (by

5 TLC) to cause varying levels of *cis*→*trans* isomerism. Finally, it was determined that trimethylsilyl iodine, neat or generated *in-situ*, provided the best route to obtain the corresponding phosphoric acid. Treatment of the latter with sodium methoxide gave, following recrystallization, a high overall yield of phosphate salt **3d**.

10 The second series of phosphorylating agents studied were the various alkylamidophosphines which have been shown to readily phosphorylate alcohols and phenols in high yield. Di-*tert*-butyloxy(N,N-diisopropylamido)phosphine **2a**, was prepared in good yield from dichloro(N,N-diisopropylamido)phosphine and *tert*-butanol. Phosphines **2b** and **2c** were prepared in a similar manner by reacting dichloro(N,N-diethylamido)phosphine and dichloro(N,N-diisopropyl-  
15 amido)phosphine with *tert*-butanol and 2-(trimethylsilyl)ethanol, respectively. In the presence of the catalyst, 1*H*-tetrazole, these reactive amides nicely phosphorylated phenol **1a** to give the corresponding phosphate esters **3b** and **3c** in greater than 90% yields. Cleavage of the ester protecting groups was easily affected with either trifluoroacetic acid for the *tert*-butyl ester or tetrabutylammonium fluoride for the *bis*(2-trimethylsilyl)ethyl ester. After isolation, the phosphoric acid derivative was treated with sodium methoxide in methanol to afford sodium phosphate **3d** in high yield.

20 The three new phosphorylation routes to prodrug **3d**, described herein, were readily employed to obtain 10 g or greater amounts without any decrease in yields or purity. The resulting phosphate prodrug **3d** samples proved to be identical with our original specimen and therefore the clinical supply requirements for combretastatin A-4 prodrug **3d** can now be met.

25 Comparison of the synthetic *trans*-stilbene phosphate **4c** (see below) with the impurity accompanying the scaleup synthesis of combretastatin A-4 prodrug **3d** confirmed the structural assignment. The ion-pair HPLC analysis noted above proved to be quite useful for detecting small quantities of this impurity. Apparently the *trans*-stilbene **4c** arose in inconsistent amounts by an

electrophilic bimolecular addition of iodine to form an iodonium ion and subsequent anti-addition product (Hassner *et al.*, 1970, Stereochemistry of halogen azide additions to olefins, The stability of three-membered iodonium *vs* bromonium ions, *Journal of the American Chemical Society*, 92, 4879; Robertson *et al.*, 1950, The kinetics of halogen addition to unsaturated compounds, Part XVIII, Iodine addition, *Journal of the Chemical Society*, 2191; Zanger *et al.*, 1975, A nuclear magnetic resonance investigation of the iodination of 1,2-disubstituted ethylenes, Evidence for a *trans* addition - *cis* elimination, *Journal of Organic Chemistry*, 40, 248; Ayres *et al.*, 1971, Reaction of iodine with olefins, Kinetics and mechanism of iodine addition to pentene isomers, *Journal of the American Chemical Society*, 93, 1389; Skell *et al.*, 1964, Stereospecific *trans* photoaddition of elementary iodine to aliphatic olefins, Bridged iodoalkyl radicals, *Journal of the American Chemical Society*, 86, 2956). Fortunately, isomeric stilbenes **3d** and **4c** have distinctive HPLC retention times in the phosphate buffer employed. The combretastatin A-4 prodrug **3d** eluted at 9.28 minutes with UV  $\lambda_{\text{max}}$  290 nm and the *trans*-isomer at 9.40 minutes with UV  $\lambda_{\text{max}}$  325 nm (see Fig 1a-1d). Because the *trans*-isomer phosphate salt **4c** proved to be sparingly soluble in water, it was simply removed from the *cis*-isomer phosphate salt **3d** by redissolving the combretastatin A-4 prodrug **3d** in a minimal amount of water and extracting with ethyl acetate. The (E)-stilbene **4c**, owing to its low solubility in water, partitioned selectively into the ethyl acetate. The water phase containing the (Z)-stilbene was concentrated and the residue recrystallized from water-acetone and methanol-acetone to give the 99+% pure (by HPLC) sodium combretastatin A-4 phosphate **3d** in >80% yields.

Since combretastatin A-4 binds tubulin very strongly in the colchicine binding site (Lin *et al.*, 1988, Interactions of tubulin with potent natural and synthetic analogs of the antimetabolic agent combretastatin, a structure-activity study, *Molecular Pharmacology*, 34, 200), with inhibition to 97% reported, it became important to replace the sodium cations in prodrug **3d** with different

5 cations, both divalent and monovalent, in order to examine their effects on activity and solubility. Using this same approach, the sodium ions were replaced with metal cations and a variety of ammonium cations in order to examine their possible effects on the cancer cell line and antimicrobial activities and solubility. Some of these ammonium cations were explored for the purpose of possibly obtaining a stable, water-soluble drug with the ability to reverse multi-drug resistance through interference with the p-glycoprotein mechanism. Watanabe *et al.*, 1997, *J. Nat. Cancer Inst.*, 89(7), pp. 512-518; Adams *et al.*, 1995 *Investigational New Drugs* 13:13-21; Sato *et al.*, 1995, *Cancer Chemother. Pharmacol.*, 35:271-277; Genne *et al.*, 1995, *Anti-cancer Drug Design*, 10, 103-118. Thus, prodrugs **5a-s** were synthesized and evaluated.

10 Once the very practical synthesis of high purity prodrug **3d** was in hand, a variety of cation derivatives of the parent phosphate were synthesized and evaluated. By treating the prodrug **3d** phosphoric acid precursor with a 1.0 M solution of the respective metal hydroxide or acetate, the corresponding salts were formed. An analogous procedure was followed for the ammonium salts, using a 1.0 M solution of the amine in methanol. For the alkaloid salts, two equivalents of the base (presuming 100% formation of the phosphoric intermediate) in a suitable solvent were added to the acid solution and allowed to react for 6-8 hours. Precipitation or recrystallization gave the products.

15 *3'-O-Bis(benzyl)phosphoryl-3,4,4',5-tetramethoxy-(E)-stilbene (4b)*

20 (E)-combretastatin A-4 (**4a**, 2.2 g, 7 mmol) was dissolved in acetonitrile (20 ml) in a flask equipped with a septa, thermometer and argon inlet. After cooling to -25°C, carbon tetrachloride (3.38 ml, 35 mmol, 5 equiv) was added and the solution stirred for 5 minutes. With a syringe, diisopropylethylamine (2.56 ml, 14.7 mmol, 2.1 equiv) was added followed by DMAP (86 mg, 0.7 mmol, 0.1 equiv). Slow (dropwise) addition of dibenzyl phosphite (2.24 ml, 10.2 mmol, 1.45 equiv) was begun one minute later at such a rate that the reaction temperature remained below -20°C. The reaction was complete in 1



hour (by TLC analysis), 0.5 M  $\text{KH}_2\text{PO}_4$  was added (5 ml), the solution allowed to warm to room temperature and extracted with ethyl acetate (3 x 20 ml). The combined solvent extract was washed with water (25 ml), saturated NaCl (25 ml) and dried. Filtration and removal of solvent gave an oil that was chromatographed on a column of silica gel (hexane-ethyl acetate, 3:2) to give phosphate ester **4b** as a clear gum (3.96 g, 98%):  $R_f$  0.15 (hexane-ethyl acetate, 3:2); EIMS  $m/z$  (% intensity) 576 ( $M^+$ , 100), 561 (12), 486 (3), 406 (6), 394 (2), 378 (6), 364 (7), 316 (17), 301 (10), 252 (8), 241 (5), 181 (5); IR (neat)  $\nu_{\text{max}}$  3005, 2939, 2839, 1734, 1581, 1512, 1456, 1425, 1346, 1278, 1126, 1008, 900, 823, 740, 698  $\text{cm}^{-1}$ ;  $^1\text{H}$ -NMR (300 MHz)  $\delta$  3.80 (s, 3H,  $\text{OCH}_3$ ), 3.87 (s, 3H,  $\text{OCH}_3$ ), 3.91 (s, 6H, 2 x  $\text{OCH}_3$ ), 5.18 (s, 2H,  $\text{CH}_2\text{-Ar}$ ), 5.21 (s, 2H,  $\text{CH}_2\text{-Ar}$ ), 6.69 (s, 2H), 6.81 (d,  $J=16.5$  Hz, 1H), 6.88 (d,  $J=16.5$  Hz, 1H), 6.89 (d,  $J=9.3$  Hz, 1H), 7.24 (m, 1H), 7.34 (s, 10H, Ar-H), 7.42 (t,  $J=3.3$  Hz, 1H);  $^{13}\text{C}$ -NMR (75 MHz)  $\delta$  153.41, 150.22, 139.87, 135.78, 133.06, 130.57, 128.53, 127.93, 127.65, 126.82, 124.28, 119.00, 112.71, 103.47, 69.95, 60.97, 56.13;  $^{31}\text{P}$ -NMR (202 MHz)  $\delta$  -5.52; Anal. calcd. for  $\text{C}_{32}\text{H}_{33}\text{O}_8\text{P}$ : C, 66.67; H, 5.77. Found: C, 66.71; H, 5.93.

*Sodium (E)-combretastatin A-4 3'-O-phosphate (4c)*

Chlorotrimethylsilane (0.88 ml, 6.9 mmol, 2 equiv) was slowly added (with vigorous stirring) to a solution of the benzyl ester (**4b**, 2.0 g, 3.5 mmol) and sodium iodide (1.04 g, 6.9 mmol, 2 equiv) in dry acetonitrile (10 ml, in a flame-dried flask under argon). After stirring for 20 minutes, TLC analysis (hexane-ethyl acetate, 3:2) showed no starting material. Enough water was added to dissolve the salts and the straw color removed by the addition of 10% aq. sodium thiosulfate (5 drops). The solvent was separated and the aqueous phase extracted with ethyl acetate (4 x 10 ml). The combined extract was concentrated *in vacuo* and the resulting foam was dissolved in dry methanol (100 ml). Sodium methoxide (95%, 375 mg, 6.9 mmol, 2 equiv) was added in one portion and the solution stirred for 6 hours. The methanol was removed under

reduced pressure and the solid recrystallized from methanol-acetone to give an off-white powder (1.43 g, 93%): m.p. 157-158°C;  $R_f$  0.45 (n-butanol-methanol-water-ammonia, 4:3:2:1); UV  $\lambda_{max}$  325 nm; IR (KBr) 2939, 2839, 1583, 1512, 1464, 1423, 1340, 1253, 1126, 991, 923, 856, 819, 773, 717, 644  $cm^{-1}$ ;  $^1H$ -NMR (500 MHz,  $CD_3OD$ )  $\delta$  3.73 (s, 3H,  $OCH_3$ ), 3.84 (s, 3H,  $OCH_3$ ), 3.87 (s, 6H, 2 x  $OCH_3$ ), 6.82 (s, 2H), 6.92 (d,  $J=8.5$  Hz, 1H), 7.00 (d,  $J=16$  Hz, 1H), 7.03 (d,  $J=16$  Hz, 1H), 7.10 (dd,  $J=8.5$  Hz, 2 Hz, 1H), 7.14 (d,  $J=2$  Hz, 1H);  $^{13}C$ -NMR (75 MHz,  $CD_3OD$ )  $\delta$  153.12, 134.07, 130.47, 127.62, 126.35, 121.20, 117.66, 112.01, 103.16, 73.70, 59.69, 55.12;  $^{31}P$ -NMR (202 MHz,  $CD_3OD$ )  $\delta$  -1.4; HRFABMS  $m/z$  441.0698  $[M + Na]^+$ , (calcd for  $C_{18}H_{20}O_8Na_2P$ , 441.0691).

#### General Procedure for Synthesis of Combretastatin A-4 Prodrugs

##### Method A (Prodrugs **5a-f**)

##### *Cesium combretastatin A-4 3'-O-phosphate (5b)*

To a stirred solution of dibenzyl combretastatin A-4 3'-O-phosphate (**4b**, 0.50 g; 0.87 mmol) and sodium iodide (0.26 g, 1.74 mmol, 2 equiv) in dry acetonitrile (4 ml) in a flame-dried flask under argon was slowly added *via* syringe chlorotrimethylsilane (0.22 ml, 1.74 mmol, 2 equiv). The mixture was stirred at room temperature until the reaction was complete by TLC (*ca* 20 min). Enough water was added to dissolve the salts and the straw color was removed by the addition of 10% sodium thiosulfate (4 drops). The aqueous phase was extracted with ethyl acetate (4 x 5 ml) and the combined extract was concentrated under reduced pressure to a light straw foam. After adding distilled water (2 ml) and 1.0 M aqueous CsOH (1.74 ml, 1.74 mmol) dropwise, the solution became cloudy. The mixture was stirred for 6 hours under argon, whereupon the precipitate was collected and recrystallized from water-acetone to afford a colorless powder (0.51 g) m.p. 167-168°C;  $^1H$ -NMR (300 MHz,  $D_2O$ )  $\delta$  3.70 (s, 6H, 2 x  $OCH_3$ ), 3.76 (s, 3H,  $OCH_3$ ), 3.83 (s, 3H,  $OCH_3$ ), 6.53 (d,  $J=12.3$  Hz, 1H), 6.65 (d,  $J=12.3$  Hz, 1H), 6.68 (s, 2H), 6.86 (d,  $J=8.4$  Hz, 1H), 6.90 (dd,  $J=8.4$ , 1.8 Hz, 1H), 7.32 (d,  $J=1.8$  Hz, 1H).

*Calcium combretastatin A-4 3'-O-phosphate (5a)*

Reprecipitated from water-acetone to give a colorless powder (147 mg): m.p. 293-295°C (dec); <sup>1</sup>H-NMR (300 MHz, D<sub>2</sub>O) δ 3.69 (s, 6H, 2 x OCH<sub>3</sub>), 3.74 (s, 3H, OCH<sub>3</sub>), 3.81 (s, 3H, OCH<sub>3</sub>), 6.54 (d, *J*=12.4 Hz, 1H), 6.67 (d, *J*=12.4 Hz, 1H), 6.69 (s, 2H), 6.85 (d, *J*=8.1 Hz, 1H), 6.91 (dd, *J*=8.1, 1.8 Hz, 1H), 7.21 (d, *J*=1.8 Hz, 1H).

*Lithium combretastatin A-4 3'-O-phosphate (5c)*

Recrystallized from water-acetone to give a colorless powder (151 mg): m.p. 241-242°C; <sup>1</sup>H-NMR (300 MHz, D<sub>2</sub>O) δ 3.56 (s, 6H, 2 x OCH<sub>3</sub>), 3.62 (s, 3H, OCH<sub>3</sub>), 3.70 (s, 3H, OCH<sub>3</sub>), 6.38 (d, *J*=12 Hz, 1H), 6.52 (s, 2H), 6.53 (d, *J*=12 Hz, 1H), 6.73 (dd, *J*=8.4, 1.4 Hz, 1H), 7.24 (bd, *J*=1.4 Hz, 1H).

*Magnesium combretastatin A-4 3'-O-phosphate (5d)*

Recrystallization from methanol-acetone afforded a creme-colored solid (121 mg): m.p. 270°C (dec); <sup>1</sup>H-NMR (300 MHz, D<sub>2</sub>O) δ 3.54 (s, 6H, 2 x OCH<sub>3</sub>), 3.62 (s, 3H, OCH<sub>3</sub>), 3.72 (s, 3H, OCH<sub>3</sub>), 6.34 (d, *J*=12.3 Hz, 1H), 6.48 (s, 2H), 6.50 (d, *J*=12.3 Hz, 1H), 6.75 (m, 2H), 7.20 (s, 1H).

*Manganese combretastatin A-4 3'-O-phosphate (5e)*

Recrystallization from water-acetone gave a creme-colored solid (143 mg): m.p. 241-242°C (dec); <sup>1</sup>H-NMR (300 MHz, D<sub>2</sub>O) δ 3.66 (s, 6H, 2 x OCH<sub>3</sub>), 3.73 (s, 3H, OCH<sub>3</sub>), 3.82 (s, 3H, OCH<sub>3</sub>), 6.46 (d, *J*=12.3 Hz, 1H), 6.58 (s, 2H), 6.59 (d, *J*=12.3 Hz, 1H), 6.87 (d, *J*=8.7 Hz, 1H), 6.94 (dd, *J*=8.7, 1.5 Hz, 1H), 7.24 (s, 1H).

*Zinc combretastatin A-4 3'-O-phosphate (5f)*

Collection of the white solid and washing with acetone gave a colorless, opalescent solid (164 mg): m.p. 198-200°C; <sup>1</sup>H-NMR (300 MHz, D<sub>2</sub>O) δ 3.58 (s, 6H, 2 x OCH<sub>3</sub>), 3.65 (s, 3H, OCH<sub>3</sub>), 3.73 (s, 3H, OCH<sub>3</sub>), 6.41 (d, *J*=12 Hz, 1H), 6.52 (s, 2H), 6.52 (d, *J*=12 Hz, 1H), 6.82 (d, *J*=8.1 Hz, 1H), 6.89 (dd, *J*=8.1, 1.4 Hz, 1H), 7.12 (d, *J*=1.4 Hz, 1H).

Method B (Prodrugs **5g-l**)

The same general procedure as described above was used, except 2 equivalents of a 1.0 M solution of the amine in dry methanol was added and the reaction mixture allowed to stir for 8 h. All the amine salts were recrystallized from methanol-ether.

*Imidazole combretastatin A-4 3'-O-phosphate (5g)*

Recrystallization gave 137 mg; m.p. 196-198°C; <sup>1</sup>H-NMR (300 MHz, D<sub>2</sub>O) δ 3.66 (s, 6H, 2 x OCH<sub>3</sub>), 3.73 (s, 3H, OCH<sub>3</sub>), 3.82 (s, 3H, OCH<sub>3</sub>), 6.47 (d, J=12 Hz, 1H), 6.58 (s, 2H), 6.59 (d, J=12 Hz, 1H), 6.92 (dd, J=8.4, 1.2 Hz, 1H), 7.23 (d, J=1.2 Hz, 1H), 7.46 (s, 2H, δ4, δ5), 8.67 (s, 1H, δ2).

*Morpholine combretastatin A-4 3'-O-phosphate (5h)*

Recrystallization afforded (121 mg 61% yield): m.p. 209-210°C; <sup>1</sup>H-NMR (300 MHz, D<sub>2</sub>O) δ 3.22 (m, 4H, -CH<sub>2</sub>OCH<sub>2</sub>-), 3.46 (s, 6H, OCH<sub>3</sub>), 3.48 (s, 3H, OCH<sub>3</sub>), 3.54 (s, 3H, OCH<sub>3</sub>), 3.88 (m, 4H, -CH<sub>2</sub>NCH<sub>2</sub>-), 6.36 (d, J=12.3 Hz, 1H), 6.49 (s, 2H), 6.50 (d, J=12.3 Hz, 1H), 6.79 (dd, J=8.1, 1.2 Hz, 1H), 7.24 (bd, 8.1 Hz, 1H), 7.40 (bd, J=1.2 Hz, 1H).

*Piperazine combretastatin A-4 3'-O-phosphate (5i)*

Following recrystallization 151 mg; m.p. 198-200°C; <sup>1</sup>H-NMR (300 MHz, CD<sub>3</sub>OD) δ 3.01 (bs, 1H, NH), 3.52 (bm, 8H, δ2-H), 3.67 (s, 6H, 2 x OCH<sub>3</sub>), 3.74 (s, 3H, OCH<sub>3</sub>), 3.83 (s, 3H, OCH<sub>3</sub>), 6.49 (d, J=12 Hz, 1H), 6.56 (d, J=12 Hz, 1H), 6.62 (s, 2H), 6.95 (m, 2H), 7.24 (s, 1H).

*Piperidine combretastatin A-4 3'-O-phosphate (5j)*

By recrystallization 137 mg (was collected); m.p. = 159-160°C; <sup>1</sup>H-NMR (300 MHz, D<sub>2</sub>O) δ 1.67 (m, 8H, [2 x δ3 H s]), 1.77 (m, 4H, [2 x δ4 H s]), 3.15 (m, 8H, [2 x δ2 H s]), 3.66 (s, 6H, 2 x OCH<sub>3</sub>), 3.73 (s, 3H, OCH<sub>3</sub>), 3.82 (s, 3H, OCH<sub>3</sub>), 6.48 (d, J=12 Hz, 1H), 6.60 (s, 2H), 6.61 (d, J=12 Hz, 1H), 6.854 (d, J=8.5 Hz, 1H), 6.92 (dd, J=19.5, 8.5 Hz, 1H), 7.26 (s, 1H).

*Pyrazole combretastatin A-4 3'-O-phosphate (5k)*

Recrystallization produced 112 mg: m.p. = 186-187°C; <sup>1</sup>H-NMR (300 MHz, D<sub>2</sub>O) δ 3.69 (s, 6H, 2 x OCH<sub>3</sub>), 3.76 (s, 3H, OCH<sub>3</sub>), 3.85 (s, 3H, OCH<sub>3</sub>), 6.30 (d, J=1.4 Hz, 2H, 2 x δ<sub>4</sub>), 6.53 (d, J=12 Hz, 1H), 6.63 (d, J=12 Hz, 1H), 6.64 (s, 2H), 6.97 (m, 2H), 7.25 (d, J=1.8 Hz, 1H), 7.67 (d, 4H, 2 x δ<sub>3</sub>, δ<sub>5</sub> H).

*Pyridine combretastatin A-4 3'-O-phosphate (5l)*

The yield was 96 mg: m.p. = 165-168°C; <sup>1</sup>H-NMR (300 MHz, D<sub>2</sub>O) δ 3.66 (s, 6H, 2 x OCH<sub>3</sub>), 3.73 (s, 3H, OCH<sub>3</sub>), 7.82 (s, 3H, OCH<sub>3</sub>), 6.48 (d, J=12 Hz, 1H), 6.58 (s, 2H), 6.59 (d, J=12 Hz, 1H), 6.90 (d, J=8.4 Hz, 1H), 6.95 (dd, J=8.4, 1.5 Hz, 1H), 7.22 (bd, J=1.5 Hz, 1H), 7.47 (d, J=3.3 Hz, 4H, 2 x δ<sub>3,5</sub>-H), 8.0 (m, 2H, δ<sub>4</sub>-H), 8.86 (m, 4H, 2 x δ<sub>2,6</sub>-H).

**Method C (Prodrugs 5m-s)**

An analogous method as summarized for prodrugs **3a-f** was used, except that two equivalents of the alkaloid or other base in the appropriate solvent were added to the phosphoric acid in the same solvent. Recrystallization/precipitation was accomplished using methanol-ether.

*Adenosine combretastatin A-4 3'-O-phosphate (5m)*

The adenosine salt weighed 160 mg: m.p. 155-158°C; <sup>1</sup>H-NMR (300 MHz, D<sub>2</sub>O) δ 3.50 (s, 6H, 2 x OCH<sub>3</sub>), 3.57 (s, 3H, OCH<sub>3</sub>), 3.68 (s, 3H, OCH<sub>3</sub>), 3.71 (d, J=2.1 Hz, 2H), 3.76 (bs, 2H), 3.80 (dd, J=2.1, 6.6 Hz, 1H), 4.14 (d, J=3.0 Hz, 2H), 4.28 (t, J=3.0 Hz, 2H), 4.58 (t, J=6.0 Hz, 2H), 5.85 (d, J=6.0 Hz, 2H), 6.21 (d, J=12.0 Hz, 1H), 6.32 (s, 2H), 6.36 (d, J=12.6 Hz, 1H), 6.71 (bs, 3H), 7.07 (s, 1H), 7.95 (d, J=1.5 Hz, 2H), 8.11 (s, 2H).

*Cinchonine combretastatin A-4 3'-O-phosphate (5n)*

The use of cinchonine led to 175 mg of product, m.p. 189-193°C; <sup>1</sup>H-NMR (300 MHz, CDCl<sub>3</sub>) δ 0.829 (m, 1H), 1.21 (m, 1H), 1.74 (m, 1H), 2.15 (t, J=12 Hz, 1H), 2.32 (q, 1H), 3.30 (br s, -OH, 1H), 3.6 (s, 3H, OCH<sub>3</sub>), 3.64 (s, 6H, 2 x OCH<sub>3</sub>), 3.78 (s, 3H, OCH<sub>3</sub>), 4.45 (t, J=9.3 Hz, 1H), 5.18 (d, J=16

Hz, 1H), 5.23 (d,  $J=10.8$  Hz, 1H), 5.42 (d,  $J=12$  Hz, 1H), 5.87 (m, 1H), 7.7 (d,  $J=8.4$  Hz, 1H), 7.76 (d,  $J=6.3$  Hz, 1H), 7.86 (d,  $J=4.5$  Hz, 1H), 7.97 (d,  $J=8.7$  Hz, 1H), 8.10 (d,  $J=8.7$  Hz, 1H), 8.29 (d,  $J=8.4$  Hz, 1H), 8.83 (d,  $J=4.5$  Hz, 1H):

5 *Glucosamine combretastatin A-4 3'-O-phosphate (5o)*

Recrystallization led to 98 mg of product, m.p. 190-195°C (dec);  $^1\text{H}$ -NMR (300 MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  1.81 (s), 1.88 (s), 1.93 (s), 2.85 (s), 2.88 (d,  $J=2.1$  Hz, 1H), 2.91 (s, 1H), 3.16 (d,  $J=3.9$  Hz, 1H), 3.19 (d,  $J=3.9$  Hz, 1H), 3.38 (h, 2H), 3.53-3.82 (m, 4H), 3.64 (s, 3H,  $\text{OCH}_3$ ), 3.69 (s, 3H,  $\text{OCH}_3$ ), 3.74 (s, 6H, 2 x  $\text{OCH}_3$ ), 4.82 (d,  $J=8.1$  Hz, 1H), 5.33 (d,  $J=3.3$  Hz, 1H), 6.51 (d,  $J=12.1$  Hz, 1H), 6.60 (d,  $J=12.1$  Hz, 1H), 6.63 (s, 2H), 6.80-7.22 (m, 3H).

10 *Quinine combretastatin A-4 3'-O-phosphate (5p)*

Recrystallization of the quinine salt gave 75 mg; m.p. 179-184°C;  $^1\text{H}$ -NMR (300 MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  3.00 (q, 10.1, 13.7 Hz, 1H), 3.15 (o,  $J=5.0$ , 8.9, 8.9 Hz, 1H), 3.37 (h,  $J=3.0$ , 5.3, 10.1, 13.2 Hz, 1H), 3.41 (bs, 1H, OH), 3.61 (s, 3H,  $\text{OCH}_3$ ), 3.69 (s, 3H,  $\text{OCH}_3$ ), 3.82 (s, 6H, 2 x  $\text{OCH}_3$ ), 3.91 (s, 3H,  $\text{OCH}_3$ ), 4.90 (d,  $J=10.0$  Hz, 1H), 4.97 (d,  $J=17.1$  Hz, 1H), 5.50 (d,  $J=4.4$  Hz, 1H), 5.84 (o,  $J=7.6$ , 10.0, 17.3 Hz, 1H), 6.55 (bd,  $J=12.6$  Hz, 1H), 6.61 (bd,  $J=12.6$  Hz, 1H), 6.63 (s, 2H), 6.67 (bs, 3H), 7.22 (s, 1H), 7.34 (d,  $J=9.5$  Hz, 1H), 7.50 (d,  $J=4.3$  Hz, 1H), 8.70 (d,  $J=4.2$  Hz, 1H).

15 *Quinidine combretastatin A-4 3'-O-phosphate (5q)*

Quinidine was used to prepare 300 mg of product, m.p. 180-181°C;  $^1\text{H}$ -NMR (300 MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  0.963 (t, 1H), 1.16 (m, 1H), 1.89 (m, 2H), 1.99 (bs, 1H), 2.34 (t,  $J=6.8$  Hz, 1H), 2.65 (dd,  $J=6.6$  Hz, 12.3 Hz, 1H), 3.10 (s, 1H), 3.22 (m, 1H), 3.29 (o,  $J=1.6$ , 7.9, 14 Hz), 3.32 (s, 3H,  $\text{OCH}_3$ ), 3.65 (s, 3H,  $\text{OCH}_3$ ), 3.99 (s, 3H,  $\text{OCH}_3$ ), 4.09 (s, 6H, 2 x  $\text{OCH}_3$ ), 4.61 (bs, 1H), 5.10 (d,  $J=9.3$  Hz), 5.28 (d,  $J=15.4$  Hz, 1H), 5.96 (bs, 1H), 6.11 (o,  $J=7.4$ , 10.5, 16.0 Hz, 1H), 6.54 (bd,  $J=12.0$  Hz, 1H), 6.60 (bd,  $J=12$  Hz, 1H), 6.61 (s, 2H), 7.43 (dd,  $J=6.9$ , 1.8 Hz, 1H), 7.56 (dd,  $J=7.2$ , 1.8 Hz, 1H), 7.61 (d,

$J=1.8$  Hz), 7.70 (t,  $J=3.3$  Hz, 1H), 7.91 (d,  $J=4.4$  Hz, 1H), 7.97 (d,  $J=9.2$  Hz, 1H), 8.06 (d,  $J=9.2$  Hz, 1H), 8.69 (d,  $J=4.4$  Hz, 1H), 8.79 (d,  $J=4.8$  Hz, 1H).

*Tetracycline combretastatin A-4 3'-O-phosphate (5r)*

The tetracycline salt upon recrystallization weighed 136 mg: m.p. 191-193°C;  $^1\text{H-NMR}$  (300 MHz,  $\text{DMSO-d}_6$ )  $\delta$  1.53-3.21 (m, 4H,  $\text{C}_4\alpha$  and  $\text{C}_5\alpha$  methine protons,  $\text{C}_5$ -methylene hydrogens), 1.56 (s, 3H,  $\text{C}_6\text{-CH}_3$ ), 2.55 (s, 6H,  $\text{N}(\text{CH}_3)_2$ ), 3.31 (bs, 1H,  $\text{C}_4\text{-H}$ ), 3.64 (s, 6H,  $2\times\text{OCH}_3$ ), 3.67 (s, 3H,  $\text{OCH}_3$ ), 3.72 (s, 3H,  $\text{OCH}_3$ ), 4.31 (bs, 1H,  $\text{C}_4\text{-H}$ ), 6.41 (d,  $J=12.6$ , 1H), 6.49 (d,  $J=12.6$ , 1H), 6.61 (s, 2H), 6.87 (s, 3H), 6.93 (d,  $J=8.1$ , 1H, D-Ar-H), 7.14 (d,  $J=7.8$ , 1H, D-Ar-H), 7.47 (s, 1H), 7.56 (t,  $J=7.8$ , 1H, D-Ar-H), 9.13 (bs, 2H,  $\text{CONH}_2$ ), 11.7 (bs, 1H,  $\text{C}_{10}\text{-OH}$ ).

*Verapamil combretastatin A-4 3'-O-phosphate (5s)*

From verapamil was obtained 110 mg of product: m.p. 185-186°C;  $^1\text{H-NMR}$  (300 MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  0.73 (d,  $J=6.6$  Hz, 3H), 1.15 (d on m,  $J=6.6$  Hz, 4H), 1.32 (m, 1H), 1.65 (m, 1H), 1.99 (s, 1H), 2.17 (m, 1H), 2.19 (s, 1H), 2.78 (bs, 2H), 3.11 (m, 2H), 3.27 (bs, 2H), 3.36 - 3.82 (4s, 8 x  $\text{OCH}_3$ , 24H), 6.47 (d,  $J=12$  Hz, 1H), 6.57 (d,  $J=12$  Hz, 1H), 6.64 (bd, 1H), 6.80 (bd, 1H), 6.92 (dd,  $J=8.4$ , 1.9 Hz, 2H), 7.28 (s, 1H).

**Biological Results**

All of the synthetic products were evaluated against a minipanel of human cancer cell lines, the murine P388 lymphocytic leukemia and a selection of microorganisms. These results are summarized in Tables I-III respectively.

In the case of Antimicrobial susceptibility testing compounds were screened against the bacteria *Stenotrophomonas maltophilia*, *Micrococcus luteus*, *Staphylococcus aureus*, *Escherichia coli*, *Enterobacter cloacae*, *Enterococcus faecalis*, *Streptococcus pneumoniae*, *Neisseria gonorrhoeae*, and the fungi *Candida albicans* and *Cryptococcus neoformans*, according to established disk susceptibility testing protocols (National Committee for Clinical Laboratory Standards, 1997).

Table I. Human Cancer Cell Line Activity ( $GI_{50}$  ( $\mu g/ml$ )) and Murine P-388 Leukemia Inhibitory Activity ( $ED_{50}$  ( $\mu g/ml$ )) of combretastatin A-4 (**1a**) and synthetic modifications (**4a-5s**).

Structure	Prostate	Pharynx	Lung-NSC	Thyroid
	DU-145	FADU	NCI-H460	SW1736
1a	0.00076	0.00065	0.00056	0.00071
3d	0.0072	0.0045	0.0035	0.0061
4a	0.037	0.040	0.038	0.080
4c	0.044	0.071	0.038	11.2
5a	0.001	0.001	0.001	0.010
5b	0.0028	0.0023	0.0032	8.4
5c	0.0025	0.003	0.004	0.030
5d	0.01	0.031	0.017	6.8
5e	<0.001	0.00091	0.0015	14.7
5f	0.0062	0.0070	0.0074	0.059
5g	0.001	0.01	0.004	0.011
5h	0.0024	0.0016	0.003	0.02
5i	0.0011	0.0018	0.0028	23.5
5j	0.01	0.01	0.01	0.50
5k	0.01	0.01	0.01	4.6
5l	0.036	0.032	0.038	0.01
5m	0.036	0.032	0.038	0.092
5n	0.030	0.024	0.037	0.089
5o	0.026	0.015	0.034	0.061
5p	0.0048	0.006	0.0042	7.5
5q	0.0036	0.0033	0.0039	0.11
5r	0.058	0.0055	0.0046	0.012
5s	0.0036	0.0046	0.004	0.0061

\* Not tested against this cell line



Table I.(continued)

Structure	Neuroblast	Pancreas	Melanoma	P-388
	SK-N-SH	BXPC-3	SK-Mel-3	
1a	0.00026	>0.1	0.00020	0.00266
3d	0.0025	0.23	NT <sup>a</sup>	0.00292
4a	0.029	3.9	NT <sup>a</sup>	0.280
4c	3.1	4.0	0.036	0.02
5a	0.015	0.22	0.0036	0.0011
5b	0.011	0.31	0.0043	0.001
5c	0.016	0.31	NT <sup>a</sup>	0.0012
5d	0.01	0.43	NT <sup>a</sup>	0.0018
5e	0.014	0.20	0.0036	0.016
5f	0.024	1.1	1.0	0.0343
5g	0.01	0.39	0.01	0.0025
5h	0.039	0.26	0.0042	0.0023
5i	0.021	0.19	0.0036	0.0016
5j	0.017	0.30	0.01	0.0029
5k	0.014	0.23	0.01	0.0023
5l	0.03	11.6	0.35	<0.001
5m	0.034	11.6	0.35	0.16
5n	NT <sup>a</sup>	1.6	NT <sup>a</sup>	0.020
5o	0.016	4.5	NT <sup>a</sup>	0.0164
5p	NT <sup>a</sup>	2.2	NT <sup>a</sup>	0.027
5q	NT <sup>a</sup>	0.43	NT <sup>a</sup>	0.0032
5r	0.0032	0.57	NT <sup>a</sup>	0.037
5s	0.0025	0.46	NT <sup>a</sup>	0.001

<sup>a</sup> Not tested against this cell line

Table II. Antimicrobial activity of combretastatin A-4 (**1a**), combretastatin A-4 Prodrug (**3d**) and synthetic modifications (**4a**, **4c**, **5a-s**).

Compound	Microbe(s) inhibited	Minimum inhibitory concentration ( $\mu\text{g}/\text{disk}$ )
1a	<i>Neisseria gonorrhoeae</i>	25-50
3d	<i>Neisseria gonorrhoeae</i>	50-100
4a	<i>Micrococcus luteus</i>	25-50
4c	<i>Neisseria gonorrhoeae</i>	50-100
5a	*	
5b	*	
5c	*	
5d	*	
5e	<i>Neisseria gonorrhoeae</i>	6.25-12.5
	<i>Candida albicans</i>	25-50
	<i>Cryptococcus neoformans</i>	12.5-25
5f	*	
5g	*	
5h	*	
5i	<i>Neisseria gonorrhoeae</i>	25-50
	<i>Candida albicans</i>	50-100
	<i>Cryptococcus neoformans</i>	25-50
5j	*	
5k	*	
5l	*	
5m	*	
5n	<i>Neisseria gonorrhoeae</i>	25-50
5o	*	
5p	*	
5q	<i>Neisseria gonorrhoeae</i>	6.25-12.5
5r	<i>Neisseria gonorrhoeae</i>	3.12-6.25
	<i>Escherichia coli</i>	3.12-6.25
	<i>Stenotrophomonas maltophilia</i>	6.25-12.5
	<i>Enterobacter cloacae</i>	3.12-6.25
	<i>Micrococcus luteus</i>	0.78-1.56
	<i>Streptococcus pneumoniae</i>	0.195-0.39
	<i>Staphylococcus aureus</i>	0.195-0.39
	<i>Enterococcus faecalis</i>	25-50
5s	*	

\*At 100  $\mu\text{g}/\text{disk}$  no inhibition of the eight bacteria and two fungi tested.

Table III. Solubility of combretastatin A-4 prodrug **3d**, (E)-stilbene prodrug **4c** and structural modifications **5a-s** in mg per 1 ml water at 25°C.

Compound	3d	4c	5a	5b	5c	5d	5e	5f	5g	5h
Solubility (mg/ml)	20	4	2	>50	6	7	3	11	10	9

Table III. (continued)

Compound	5i	5j	5k	5l	5m	5n	5o	5p	5q	5r	5s
Solubility (mg/ml)	1	5	8	3	4	4	15	6	2	1	16

Combretastatin A-4, combretastatin A-4 prodrug and several prodrug modifications inhibited growth of the pathogenic bacterium *Neisseria gonorrhoeae* (Table II). Addition of manganese **5e** and piperazine **5i** to the parent phosphate yielded compounds inhibitory to opportunistic fungi. As expected, the tetracycline derivative **5r** inhibited the growth of all Gram-negative and Gram-positive bacteria tested.

### Biological Evaluation and Statistical Definitions

The following measures are used to express drug activity by giving the drug dose which reduces cell growth to a specified percentage of growth:

**ED<sub>50</sub>** (P388) and **GI<sub>50</sub>** (HTCL) are the drug doses (in  $\mu\text{g/ml}$  of cancer cells) needed to reduce the cancer cells by 50%. There is no mathematical difference between **ED<sub>50</sub>** and **GI<sub>50</sub>**, which are both calculated using the same formula. The only difference is historical usage.

Total Growth Inhibition ("TGI") is the drug dose (in  $\mu\text{g/ml}$  of cancer cells) needed to yield zero percent growth. Whether just as many cells were killed as were produced (steady state), or no growth occurred (total inhibition), cannot be distinguished.

Lethal Concentration 50% ("LC<sub>50</sub>") is the drug concentration (in  $\mu\text{g/ml}$  of cancer cells) kills half of the cells originally present at the beginning of the experiment.

Each drug is tested at five (5) doses: 100 - 10 - 1 - 0.1 - 0.01  $\mu\text{g/ml}$ . Percent Growths are calculated for each dose. The two (or three) doses with growth values above, below, (or near to) 50% growth are used to calculate the ED<sub>50</sub>/GI<sub>50</sub> using a linear regression formula. The log of the dose is used during the regression computation. If no dose yields a growth value under 50%, the results are expressed as: ED<sub>50</sub> > (highest dose). If no dose yields growth higher than 50% growth, then ED<sub>50</sub> < (lowest dose). Similar calculations are performed for the TGI at 0% growth, and at -50% growth for the LC<sub>50</sub>.

#### PERCENT OF GROWTH

At the start of an experiment, cells from the in vitro cell cultures are inoculated into the appropriate tubes or microtiter plates. One set of control tubes/plates is immediately counted to determine the number of cells at the start of the experiment. This is the "baseline count", or "T<sub>zero</sub> reading". At the end of the experiment (48 hours later) a second set of control tubes/plates is analyzed to determine the "Control Growth" value. The growth (or death) of cells relative to the initial quantity of cells is used to define the "Percent of Growth".

## EXAMPLE:

Baseline Count = 20

Control Count = 200

(10-Fold Growth)

100% Growth = Control Growth

100% Growth = 200

$$50\% \text{ Growth} = T_{\text{zero}} + \frac{\text{Control} - T_{\text{zero}}}{2}$$

50% Growth = 110

0% Growth =  $T_{\text{zero}}$ 

0% Growth = 20

-50% Growth =  $T_{\text{zero}} / 2$ 

-50% Growth = 10

For further information about the testing protocols and procedures see Anne Monks et al., "Feasibility of a High-Flux Anticancer Drug Screen Using a Diverse Panel of Cultured Human Tumor Cell Lines", 83 J. Nat. Cancer Inst., No. 11, pp. 757-66 (5 June 1991) and Michael J. Boyd, "Status of the NCI Preclinical Antitumor Drug Discovery Screen", 3 Princ. & Practice of Oncology Updates, No. 10, pp. 1-12 (Oct. 1989).

By definition a "prodrug" is a precursor which will undergo metabolic activation in vivo to the active drug. Thus, certain phosphate derivatives make ideal prodrugs if the phosphate group can be cleaved by endogenous non-specific phosphatases (McComb *et al.*, Alkaline Phosphatase, New York: Plenum Press, 1979) As previously reported (*See*, U.S. Patent Nos. 4,996,237 and 5,561,122), a preliminary in vitro comparison of the stable combretastatin A-4 phosphate metal or ammonium salts against the P388 murine leukemia and a selection of six human tumor cell lines (OVCAR-3, SF-295, A498, NCI-H460, KM20L2, SK-MEL-5) was performed using the sulforhodamine B assay (Skehan *et al.*, "New Colorimetric Cytotoxicity Assay for Anticancer-Drug Screening," J. Nat. Cancer Inst., 1107 (1990)). Again, sodium phosphate derivative **3d** is an equivalent to the derivative **1h** disclosed earlier. The assay results relative to **1h** will thus be repeated here with respect to **3d**. Combretastatin A-4 **1a** and prodrug **1h** (**3d**) were then comparatively evaluated against the full-panel NCI

screen (Boyd, "Status of the NCI Preclinical Antitumor Drug Discovery Screen: Implications for Selection of New Agents for Clinical trial," 3 Cancer: Principles and Practice of Oncology Updates, No. 10, pp. 1-12 (1989); and Boyd et al., "The Future of New Drug Development, Section I, Introduction to Cancer Therapy," Current Therapy in Oncology, Philadelphia: Decker (1993)). Both were tested in quadruplicate at each of three different concentration ranges ( $10^{-5}$ ,  $10^{-6}$  and  $10^{-7}$ M upper limits, five,  $\log_{10}$ -spaced concentrations in each range). Optimal tests were selected for overall potency and differential cytotoxicity comparisons (Boyd and Paul, "Some Practical Considerations and Applications of the NCI in vitro Anticancer Drug Discovery Screen," Drug Development Research, (In Press)). The mean panel  $GI_{50}$  concentration for the parent compound **1a** was  $6.61 \pm 0.79 \times 10^{-9}$ M, compared to  $6.89 \pm 0.96 \times 10^{-9}$ M for prodrug **1h (3d)**. TGI-COMPARE analyses (Id.) revealed a very high Pearson correlation coefficient of 0.91 for the differential cytotoxicity profiles of combretastatin A-4 **1a** and its prodrug **1h (3d)**. This high correlation coefficient reflects similarities in biological properties and/or chemical structure and properties. Paull *et al.*, "Display and analysis of Patterns of Differential Activity of Drugs Against Human Tumor Cell Lines: Development of Mean Graph and COMPARE Algorithm," 81 J. Nat. Cancer Inst., No. 14, pp. 108-92 (July 19, 1989).

Clearly, the excellent water solubility of sodium phosphate **1h (3d)**, good stability and cell growth inhibitory activity comparable to combretastatin **1a** allowed selection of this prodrug candidate for drug formulation studies. In a like manner, it is believed that any of the combretastatin A-4 phosphate metal or ammonium salts can be synthesized and utilized as prodrugs with cell growth inhibitory activity comparable to combretastatin **1a**.

Based upon the foregoing, compositions **1h (3d)** are believed useful in the treatment of one or more neoplastic diseases. For example, acute myelocytic leukemia, acute lymphocytic leukemia, malignant melanoma, adenocarcinoma of lung, neuroblastoma, small cell carcinoma of the lung, breast carcinoma, colon carcinoma, ovarian carcinoma, bladder carcinoma, and the like.

The dosage administered will be dependent upon the identity of the neoplastic disease; the type of host involved, including its age, health and

weight; the kind of concurrent treatment, if any; the frequency of treatment and therapeutic ratio.

Illustratively, dosage levels of the administered active ingredients are: intravenous, 0.1 to about 20 mg/kg; intramuscular, 1 to about 50 mg/kg; orally, 5 to about 100 mg/kg; intranasal instillation, 5 to about 100 mg/kg; and aerosol, 5 to about 100 mg/kg of host body weight.

Expressed in terms of concentration, an active ingredient can be present in the compositions of the present invention for localized use about the cutis, intranasally, pharyngolaryngeally, bronchially, intravaginally, rectally, or ocularly in a concentration of from about 0.01 to about 50% w/w of the composition; preferably about 1 to about 20% w/w of the composition; and for parenteral use in a concentration of from about 0.05 to about 50% w/v of the composition and preferably from about 5 to about 20% w/v.

The compositions of the present invention are preferably presented for administration to humans and animals in unit dosage forms, such as tablets, capsules, pills, powders, granules, suppositories, sterile parenteral solutions or suspensions, sterile non-parenteral solutions of suspensions, and oral solutions or suspensions and the like, containing suitable quantities of an active ingredient.

For oral administration either solid or fluid unit dosage forms can be prepared.

Powders are prepared quite simply by comminuting the active ingredient to a suitably fine size and mixing with a similarly comminuted diluent. The diluent can be an edible carbohydrate material such as lactose or starch. Advantageously, a sweetening agent or sugar is present as well as a flavoring oil.

Capsules are produced by preparing a powder mixture as hereinbefore described and filling into formed gelatin sheaths. Advantageously, as an adjuvant to the filling operation, a lubricant such as talc, magnesium stearate, calcium stearate and the like is added to the powder mixture before the filling operation.

Soft gelatin capsules are prepared by machine encapsulation of a slurry of active ingredients with an acceptable vegetable oil, light liquid petrolatum or other inert oil or triglyceride.

5 Tablets are made by preparing a powder mixture, granulating or slugging, adding a lubricant and pressing into tablets. The powder mixture is prepared by mixing an active ingredient, suitably comminuted, with a diluent or base such as starch, lactose, kaolin, dicalcium phosphate and the like. The powder mixture can be granulated by wetting with a binder such as corn syrup, gelatin solution, methylcellulose solution or acacia mucilage and forcing through a screen. As an alternative to granulating, the powder mixture can be slugged, i.e., run through the tablet machine and the resulting imperfectly formed tablets broken into pieces (slugs). The slugs can be lubricated to prevent sticking to the tablet-forming dies by means of the addition of stearic acid, a stearic salt, talc or mineral oil. The lubricated mixture is then compressed into tablets.

10 Advantageously, the tablet can be provided with a protective coating consisting of a sealing coat or enteric coat of shellac, a coating of sugar and methylcellulose and polish coating of carnauba wax.

15 Fluid unit dosage forms for oral administration such as in syrups, elixirs and suspensions can be prepared wherein each teaspoonful of composition contains a predetermined amount of an active ingredient for administration. The water-soluble forms can be dissolved in an aqueous vehicle together with sugar, flavoring agents and preservatives to form a syrup. An elixir is prepared by using a hydroalcoholic vehicle with suitable sweeteners together with a flavoring agent. Suspensions can be prepared of the insoluble forms with a suitable vehicle with the aid of a suspending agent such as acacia, tragacanth, methylcellulose and the like.

20 For parenteral administration, fluid unit dosage forms are prepared utilizing an active ingredient and a sterile vehicle, water being preferred. The active ingredient, depending on the form and concentration used, can be either suspended or dissolved in the vehicle. In preparing solutions the water-soluble active ingredient can be dissolved in water for injection and filter sterilized before filling into a suitable vial or ampule and sealing. Advantageously, adjuvants such as a local anesthetic, preservative and buffering agents can be dissolved in the vehicle. Parenteral suspensions are prepared in substantially the same manner except that an active ingredient is suspended in the vehicle



5 instead of being dissolved and sterilization cannot be accomplished by filtration. The active ingredient can be sterilized by exposure to ethylene oxide before suspending in the sterile vehicle. Advantageously, a surfactant or wetting agent is included in the composition to facilitate uniform distribution of the active ingredient.

10 In addition to oral and parenteral administration, the rectal and vaginal routes can be utilized. An active ingredient can be administered by means of a suppository. A vehicle which has a melting point at about body temperature or one that is readily soluble can be utilized. For example, cocoa butter and various polyethylene glycols (Carbowaxes) can serve as the vehicle.

15 For intranasal instillation, a fluid unit dosage form is prepared utilizing an active ingredient and a suitable pharmaceutical vehicle, preferably P.F. water, a dry powder can be formulated when insufflation is the administration of choice.

20 For use as aerosols, the active ingredients can be packaged in a pressurized aerosol container together with a gaseous or liquified propellant, for example, dichlorodifluoromethane, carbon dioxide, nitrogen, propane, and the like, with the usual adjuvants such as cosolvents and wetting agents, as may be necessary or desirable.

25 The term "unit dosage form" as used in the specification and claims refers to physically discrete units suitable as unitary dosages for human and animal subjects, each unit containing a predetermined quantity of active material calculated to produce the desired therapeutic effect in association with the required pharmaceutical diluent, carrier or vehicle. The specifications for the novel unit dosage forms of this invention are dictated by and are directly dependent on (a) the unique characteristics of the active material and the particular therapeutic effect to be achieved, and (b) the limitation inherent in the art of compounding such an active material for therapeutic use in humans, as disclosed in this specification, these being features of the present invention. 30 Examples of suitable unit dosage forms in accord with this invention are tablets, capsules, troches, suppositories, powder packets, wafers, cachets, teaspoonfuls, tablespoonfuls, dropperfuls, ampules, vials, segregated multiples of any of the foregoing, and other forms as herein described.

5 The active ingredients to be employed as antineoplastic agents can be easily prepared in such unit dosage form with the employment of pharmaceutical materials which themselves are available in the art and can be prepared by established procedures. The following preparations are illustrative of the preparation of the unit dosage forms of the present invention, and not as a limitation thereof. Several dosage forms were prepared embodying the present invention. They are shown in the following examples in which the notation "active ingredient" signifies either of the combretastatin A-4 phosphate metal or ammonium salts, e.g. **3d**, or structural modifications **4a**, **4c** and/or **5a-s**.

#### COMPOSITION "A"

##### Hard-Gelatin Capsules

15 One thousand two-piece hard gelatin capsules for oral use, each capsule containing 200 mg of an active ingredient are prepared from the following types and amounts of ingredients:

Active ingredient, micronized	200 g
Corn Starch	20 g
Talc	20 g
Magnesium stearate	2 g

20 The active ingredient, finely divided by means of an air micronizer, is added to the other finely powdered ingredients, mixed thoroughly and then encapsulated in the usual manner.

25 The foregoing capsules are useful for treating a neoplastic disease by the oral administration of one or two capsules one to four times a day.

Using the procedure above, capsules are similarly prepared containing an active ingredient in 50, 250 and 500 mg amounts by substituting 50 g, 250 g and 500 g of an active ingredient for the 200 g used above.

#### COMPOSITION "B"

##### Soft Gelatin Capsules

30 One-piece soft gelatin capsules for oral use, each containing 200 mg of an active ingredient, finely divided by means of an air micronizer, are prepared

by first suspending the compound in 0.5 ml of corn oil to render the material capsulatable and then encapsulating in the above manner.

The foregoing capsules are useful for treating a neoplastic disease by the oral administration of one or two capsules one to four times a day.

### COMPOSITION "C"

#### Tablets

One thousand tablets, each containing 200 mg of an active ingredient, are prepared from the following types and amounts of ingredients:

Active ingredient, micronized	200 g
Lactose	300 g
Corn starch	50 g
Magnesium stearate	4 g
Light liquid petrolatum	5 g

The active ingredient, finely divided by means of an air micronizer, is added to the other ingredients and then thoroughly mixed and slugged. The slugs are broken down by forcing them through a Number Sixteen screen. The resulting granules are then compressed into tablets, each tablet containing 200 mg of the active ingredient.

The foregoing tablets are useful for treating a neoplastic disease by the oral administration of one or two tablets one to four times a day.

Using the procedure above, tablets are similarly prepared containing an active ingredient in 250 mg and 100 mg amounts by substituting 250 g and 100 g of an active ingredient for the 200 g used above.

### COMPOSITION "D"

#### Oral Suspension

One liter of an aqueous suspension for oral use, containing in each teaspoonful (5 ml) dose, 50 mg of an active ingredient, is prepared from the following types and amounts of ingredients:

Active ingredient, micronized	10 g
Citric acid	2 g
Benzoic acid	1 g
Sucrose	790 g
Tragacanth	5 g

Lemon Oil 2 g  
Deionized water, q.s. 1000 ml

The citric acid, benzoic acid, sucrose, tragacanth and lemon oil are dispersed in sufficient water to make 850 ml of suspension. The active ingredient, finely divided by means of an air micronizer, is stirred into the syrup unit uniformly distributed. Sufficient water is added to make 1000 ml.

The composition so prepared is useful for treating a neoplastic disease at a dose of 1 teaspoonful (15 ml) three times a day.

#### COMPOSITION "E"

##### Parenteral Product

A sterile aqueous suspension for parenteral injection, containing 30 mg of an active ingredient in each milliliter for treating a neoplastic disease, is prepared from the following types and amounts of ingredients:

Active ingredient, micronized	30 g
POLYSORBATE 80	5 g
Methylparaben	2.5 g
Propylparaben	0.17 g
Water for injection, q.s.	1000 ml.

All the ingredients, except the active ingredient, are dissolved in the water and the solution sterilized by filtration. To the sterile solution is added the sterilized active ingredient, finely divided by means of an air micronizer, and the final suspension is filled into sterile vials and the vials sealed.

The composition so prepared is useful for treating a neoplastic disease at a dose of 1 milliliter (1ml) three times a day.

#### COMPOSITION "F"

##### Suppository, Rectal and Vaginal

One thousand suppositories, each weighing 2.5 g and containing 200 mg of an active ingredient are prepared from the following types and amounts of ingredients:

Active ingredient, micronized	15 g
Propylene glycol	150 g
Polyethylene glycol #4000, q.s.	2,500 g

5 The active ingredient is finely divided by means of an air micronizer and added to the propylene glycol and the mixture passed through a colloid mill until uniformly dispersed. The polyethylene glycol is melted and the propylene glycol dispersion is added slowly with stirring. The suspension is poured into unchilled molds at 40°C. The composition is allowed to cool and solidify and then removed from the mold and each suppository foil wrapped.

The foregoing suppositories are inserted rectally or vaginally for treating a neoplastic disease.

#### COMPOSITION "G"

##### Intranasal Suspension

10 One liter of a sterile aqueous suspension for intranasal instillation, containing 20 mg of an active ingredient in each milliliter, is prepared from the following types and amounts of ingredients:

Active ingredient, micronized	15 g
POLYSORBATE 80	5 g
Methylparaben	2.5 g
Propylparaben	0.17 g
Deionized water, q.s.	1000 ml.

15 All the ingredients, except the active ingredient, are dissolved in the water and the solution sterilized by filtration. To the sterile solution is added the sterilized active ingredient, finely divided by means of an air micronizer, and the final suspension is aseptically filled into sterile containers.

20 The composition so prepared is useful for treating a neoplastic disease, by intranasal instillation of 0.2 to 0.5 ml given one to four times per day.

25 An active ingredient can also be present in the undiluted pure form for use locally about the cutis, intranasally, pharyngolaryngeally, bronchially, or orally.

#### COMPOSITION "H"

##### Powder

30 Five grams of an active ingredient in bulk form is finely divided by means of an air micronizer. The micronized powder is placed in a shaker-type container.

The foregoing composition is useful for treating a neoplastic disease, at localized sites by applying a powder one to four times per day.

#### COMPOSITION "I"

##### Oral Powder

One hundred grams of an active ingredient in bulk form is finely divided by means of an air micronizer. The micronized powder is divided into individual doses of 200 mg and packaged.

The foregoing powders are useful for treating a neoplastic disease, by the oral administration of one or two powders suspended in a glass of water, one to four times per day.

#### COMPOSITION "J"

##### Insufflation

One hundred grams of an active ingredient in bulk form is finely divided by means of an air micronizer.

The foregoing composition is useful for treating a neoplastic disease, by the inhalation of 300 mg one to four times a day.

\* \* \*

From the foregoing, it becomes readily apparent that new and useful antineoplastic drugs and new and useful antineoplastic preparations based thereupon have been herein described and illustrated which fulfill all of the aforesaid objectives in a remarkably unexpected fashion. It is of course understood that such modifications, alterations and adaptations as will readily occur to the artisan confronted with this disclosure are intended within the spirit of the present invention as defined by the following claims.

## Claims

1. The method of synthesizing combretastatin A-4 prodrugs and trans-combretastatin A-4 prodrugs comprising:

5           admixing combretastatin A-4 with a phosphorylating agent to form a phosphate ester of combretastatin A-4 having protective groups thereupon;

10           selectively cleaving said phosphate ester protective groups with iodotrimethylsilane to yield a phosphoric acid derivative of combretastatin A-4; and

15           treating said phosphoric acid derivative of combretastatin A-4 with sodium methoxide to yield combretastatin A-4 Prodrug or a trans-isomer thereof as the ultimate product.

20           2. The method according to Claim 1 in which said phosphorlating agent is selected from the group consisting of dibenzyl phosphite/carbon tetrachloride, di-tert-butyloxy (N,N-diethylamido) phosphine and bis[2-(trimethylsilyl)ethoxy]N,N-diisopropylamidophosphine.

25           3. The method of Claim 1 in which said cleaving agent is selected from the group consisting of: iodotrimethylsilane, trifluoroacetic acid, and tetrabutylammonium fluoride.

30           4. The method of Claim 2 in which said cleaving agent is selected from the group consisting of: iodotrimethylsilane, trifluoroacetic acid, and tetrabutylammonium fluoride.

            5. The method of claim 1 in which the ultimate product consists of X-combretastatin A-4 3'-O-phosphate wherein "X" is selected from the group consisting of cesium, calcium, lithium, magnesium, manganese and zinc.

6. The method of claim 1 in which the ultimate product consists of Y-combretastatin A-4 3'-O phosphate wherein "Y" is selected from the group consisting of imidazole, morpholine, piperazine, piperidine, pyrazole and pyridine.

7. The method of claim 1 in which the ultimate product is Z-combretastatin A-4 3'-O phosphate wherein "Z" is selected from the group consisting of adenosine, cinchonine, glucosamine, quinine, quinidine, tetracycline, and verapamil.

8. The method of synthesizing a trans-isomer of combretastatin A-4 Pro-drug comprising:

dissolving combretastatin A-4 in acetonitrile;

cooling the solution to -25°F;

adding carbon tetrachloride to the cooled solution with stirring;

adding 4-dimethylaminopyridine and dibenzyl phosphate to the cooled, stirred solution and warm to room temperature;

extracting the solvent from said room temperature solution to provide 3'-O-Bis(benzyl) phosphoryl- 3, 4, 4', 5 - tetramethoxy-(E)-stilbene;

admixing chlorotrimethyl silane with said 3'-O-Bis(benzyl) phosphoryl- 3, 4, 4', 5 - tetramethoxy-(E) stilbene;

separating the solvent from the admixed solution with ethyl acetate to form an extract;


dissolving the extract in methanol;



adding sodium methoxide to said dissolved extract to form a second solution; and

removing the methanol from the second solution and recrystallizing the solid from said second solution to form a trans-isomer of combretastatin A-4 prodrug.

9. Combretastatin A-4 metal and ammonium phosphate prodrugs, and trans-combretastatin A-4 prodrugs having the structure



wherein Y is selected from the group consisting of cesium, calcium, lithium, magnesium, manganese, zinc, imidazole, morpholine, piperazine, piperidine, pyrazole, pyridine, adenosine, cinchonine, glucosamine, quinine, quinidine, tetracycline, and verapamil.

## AMENDED CLAIMS

[received by the International Bureau on 24 May 1999 (24.05.99);  
original claims 1-5 and 8-9 amended; new claim 10 added  
remaining claims unchanged (4 pages)]

1. The method of synthesizing combretastatin A-4 prodrugs as  
monosodium and disodium phosphate salts and trans-combretastatin A-4  
prodrugs comprising:

admixing combretastatin A-4 with a phosphorylating agent to form a  
phosphate ester of combretastatin A-4 having protective groups thereupon;

selectively cleaving said phosphate ester protective groups with a  
cleaving agent to yield a phosphoric acid derivative of combretastatin A-4; and

treating said phosphoric acid derivative of combretastatin A-4 with  
sodium methoxide to yield combretastatin A-4 prodrug disodium phosphate, the  
combretastatin A-4 prodrug monosodium phosphate or a trans-isomer thereof as  
the ultimate product.

2. The method according to Claim 1 in which said phosphorylating  
agent is selected from the group consisting of dibenzyl phosphite/carbon  
tetrachloride, di-tert-butoxy (N,N-diethylamido) phosphine and bis[2-  
(trimethylsilyl)ethoxy]N,N-diisopropylamidophosphine.

3. The method of Claim 1 in which said cleaving agent is selected from  
the group consisting of: iodotrimethylsilane, trifluoroacetic acid, and  
tetrabutylammonium fluoride.

4. The method of Claim 2 in which said cleaving agent is selected from  
the group consisting of: iodotrimethylsilane, trifluoroacetic acid, and  
tetrabutylammonium fluoride.

5 5. The method of claim 1 in which the ultimate product consists of X-combretastatin A-4 3'-O-phosphate wherein "X" is selected from the group consisting of cesium, calcium, lithium, magnesium, manganese, sodium, potassium and zinc.

10 6. The method of claim 1 in which the ultimate product consists of Y-combretastatin A-4 3'-O phosphate wherein "Y" is selected from the group consisting of imidazole, morpholine, piperazine, piperidine, pyrazole and pyridine.

15 7. The method of claim 1 in which the ultimate product is Z-combretastatin A-4 3'-O phosphate wherein "Z" is selected from the group consisting of adenosine, cinchonine, glucosamine, quinine, quinidine, tetracycline, and verapamil.

20 8. The method of synthesizing a trans-isomer of combretastatin A-4 Pro-drug comprising:

dissolving combretastatin A-4 in acetonitrile;

25 cooling the solution to -25°F;

adding carbon tetrachloride to the cooled solution with stirring;

30 adding 4-dimethylaminopyridine and dibenzyl phosphate to the cooled, stirred solution and warm to room temperature;

extracting the solvent from said room temperature solution to provide 3'-O-Bis(benzyl) phosphoryl- 3, 4, 4', 5 - tetramethoxy-(E)-stilbene;

admixing chlorotrimethyl silane with said 3'-O-Bis(benzyl) phosphoryl-  
3, 4, 4', 5 - tetramethoxy-(E) stilbene;


separating the solvent from the admixed solution with ethyl acetate to  
form an extract;

dissolving the extract in methanol;

adding sodium methoxide to said dissolved extract to form a second  
solution; and

removing the methanol from the second solution and recrystallizing the  
solid from said second solution to form a trans-isomer of combretastatin A-4  
prodrug.

9. Combretastatin A-4 metal and ammonium phosphate prodrugs, and  
trans-combretastatin A-4 prodrugs having the structure



wherein X is selected from the group consisting of cesium, calcium, lithium,  
magnesium, manganese, sodium, potassium, zinc, imidazole, morpholine,  
piperazine, piperidine, pyrazole, pyridine, adenosine, cinchonine, glucosamine,  
quinine, quinidine, tetracycline, and verapamil.

10. The method of claim 1 in which the ultimate product is recrystallized.

a




FIGURE 1

2/2




FIGURE 1 (continued)

Please type a plus sign (+) inside this box → ☐

PTO/SB/01 (12-97)

Approved for use through 9/30/00. OMB 0651-0032

Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it contains a valid OMB control number.

**DECLARATION FOR UTILITY OR  
DESIGN  
PATENT APPLICATION  
(37 CFR 1.63)**

☐ Declaration Submitted with Initial Filing **OR** ☐ Declaration Submitted after Initial Filing (surcharge (37 CFR 1.16 (e)) required)

Attorney Docket Number 5368-US

First Named Inventor G.R. PETITT

**COMPLETE IF KNOWN**

Application Number /

Filing Date

Group Art Unit

Examiner Name

As a below named inventor, I hereby declare that:

My residence, post office address, and citizenship are as stated below next to my name.

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

SYNTHESIS OF COMBRETASTATIN A-4 PRODRUGS AND TRANS-ISOMERS THEREOF

the specification of which (Title of the Invention)

☐ is attached hereto  
OR

☒ was filed on (MM/DD/YYYY) 01/08/1999 as United States Application Number or PCT International

Application Number PCT/US99/00419 and was amended on (MM/DD/YYYY) (if applicable).

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment specifically referred to above.

I acknowledge the duty to disclose information which is material to patentability as defined in 37 CFR 1.56.

I hereby claim foreign priority benefits under 35 U.S.C. 119(a)-(d) or 365(b) of any foreign application(s) for patent or inventor's certificate, or 365(a) of any PCT international application which designated at least one country other than the United States of America, listed below and have also identified below, by checking the box, any foreign application for patent or inventor's certificate, or of any PCT international application having a filing date before that of the application on which priority is claimed.

Prior Foreign Application Number(s)	Country	Foreign Filing Date (MM/DD/YYYY)	Priority Not Claimed	Certified Copy Attached?	
				YES	NO
			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

☐ Additional foreign application numbers are listed on a supplemental priority data sheet PTO/SB/02B attached hereto:

I hereby claim the benefit under 35 U.S.C. 119(e) of any United States provisional application(s) listed below.

Application Number(s)	Filing Date (MM/DD/YYYY)	<input type="checkbox"/> Additional provisional application numbers are listed on a supplemental priority data sheet PTO/SB/02B attached hereto.
60/071,070	01/09/1998	
60/111,531	12/09/1998	

[Page 1 of 2]

Burden Hour Statement: This form is estimated to take 0.4 hours to complete. Time will vary depending upon the needs of the individual case. Any comments on the amount of time you are required to complete this form should be sent to the Chief Information Officer, Patent and Trademark Office, Washington, DC 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Assistant Commissioner for Patents, Washington, DC 20231.



Please type a plus sign (+) inside this box → ☐

PTO/SB/01 (12-97)  
Approved for use through 9/30/00. OMB 0651-0032  
Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE  
Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it contains a valid OMB control number.

## DECLARATION — Utility or Design Patent Application

I hereby claim the benefit under 35 U.S.C. 120 of any United States application(s), or 365(c) of any PCT international application designating the United States of America, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT International application in the manner provided by the first paragraph of 35 U.S.C. 112, I acknowledge the duty to disclose information which is material to patentability as defined in 37 CFR 1.56 which became available between the filing date of the prior application and the national or PCT international filing date of this application.

U.S. Parent Application or PCT Parent Number	Parent Filing Date (MM/DD/YYYY)	Parent Patent Number (if applicable)

☐ Additional U.S. or PCT international application numbers are listed on a supplemental priority data sheet PTO/SB/02B attached hereto.

As a named inventor, I hereby appoint the following registered practitioner(s) to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith:

☐ Customer Number

OR

☒ Registered practitioner(s) name/registration number listed below

Place Customer  
Number Bar Code  
Label here

Name	Registration Number	Name	Registration Number
Richard R. Mybeck	17,886		

☐ Additional registered practitioner(s) named on supplemental Registered Practitioner Information sheet PTO/SB/02C attached hereto.

Direct all correspondence to: ☐ Customer Number  OR ☒ Correspondence address below

Name	Richard R. Mybeck				
Address	8010 East Morgan Trail, Suite 10				
Address					
City	Scottsdale	State	AZ	ZIP	85258
Country	US	Telephone	480.483.1285	Fax	480.483.7452

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under 18 U.S.C. 1001 and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Name of Sole or First Inventor: ☐ A petition has been filed for this unsigned inventor

Given Name (first and middle (if any))	Family Name or Surname
GEORGE R.	PETTIT

Inventor's Signature	<i>George R. Pettit</i>	Date	6/26/00				
Residence: City	Paradise Valley	State	AZ	Country	United States	Citizenship	US
Post Office Address	6232 Bret Hills Drive						
Post Office Address							
City	Paradise Valley	State	Arizona	ZIP	85253	Country	US

☒ Additional inventors are being named on the 1 supplemental Additional Inventor(s) sheet(s) PTO/SB/02A attached hereto

Please type a plus sign (+) inside this box → +

PTO/SB/02A (3-97)  
Approved for use through 9/30/98. OMB 0651-0032  
Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it contains a valid OMB control number.

## DECLARATION

ADDITIONAL INVENTOR(S)  
Supplemental Sheet  
Page 1 of 1

Name of Additional Joint Inventor, if any:

☐ A petition has been filed for this unsigned inventor

Given Name (first and middle [if any])

Family Name or Surname

MONTE R.

RHODES

Inventor's  
Signature

*Monte R. Rhodes*

Date

7/1/2000

Residence: City

Marcellus

State

MI

Country

United States

Citizenship

US

Post Office Address

10526 Cranberry Lake Road

Post Office Address

City

Marcellus

State

MI

ZIP

49067

Country

United States

Name of Additional Joint Inventor, if any:

☐ A petition has been filed for this unsigned inventor

Given Name (first and middle [if any])

Family Name or Surname

Inventor's  
Signature

Date

Residence: City

State

Country

Citizenship

Post Office Address

Post Office Address

City

State

ZIP

Country

Name of Additional Joint Inventor, if any:

☐ A petition has been filed for this unsigned inventor

Given Name (first and middle [if any])

Family Name or Surname

Inventor's  
Signature

Date

Residence: City

State

Country

Citizenship

Post Office Address

Post Office Address

City

State

ZIP

Country

Burden Hour Statement: This form is estimated to take 0.4 hours to complete. Time will vary depending upon the needs of the individual case. Any comments on the amount of time you are required to complete this form should be sent to the Chief Information Officer, Patent and Trademark Office, Washington, DC 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Assistant Commissioner for Patents, Washington, DC 20231.